

PTEN (N-19): sc-6818

BACKGROUND

As human tumors progress to advanced stages, one genetic alteration that occurs at high frequency is a loss of heterozygosity (LOH) at chromosome 10q23.31. Mapping of homozygous deletions on this chromosome led to the isolation of the PTEN gene, also designated MMAC1 (for mutated in multiple advanced cancers) and TEP1. This candidate tumor suppressor gene exhibits a high frequency of mutations in human glioblastomas and is also mutated in other cancers, including sporadic brain, breast, kidney and prostate cancers. PTEN has been associated with Cowden disease, an autosomal dominant cancer predisposition syndrome. The PTEN gene product is a putative protein tyrosine phosphatase that is localized to the cytoplasm, and it shares extensive homology with the cytoskeletal proteins tensin and auxilin. Gene transfer studies have indicated that the phosphatase domain of PTEN is essential for growth suppression of glioma cells.

CHROMOSOMAL LOCATION

Genetic locus: PTEN (human) mapping to 10q23.31; Pten (mouse) mapping to 19 C1.

SOURCE

PTEN (N-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of PTEN of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-6818 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

PTEN (N-19) is recommended for detection of PTEN of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

PTEN (N-19) is also recommended for detection of PTEN in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for PTEN siRNA (h): sc-29459, PTEN siRNA (m): sc-36326, PTEN shRNA Plasmid (h): sc-29459-SH, PTEN shRNA Plasmid (m): sc-36326-SH, PTEN shRNA (h) Lentiviral Particles: sc-29459-V and PTEN shRNA (m) Lentiviral Particles: sc-36326-V.

Molecular Weight of PTEN: 55 kDa.

Positive Controls: A-431 whole cell lysate: sc-2201, KNRK whole cell lysate: sc-2214 or PTEN (m): 293T Lysate: sc-122834.


STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.


RESEARCH USE

For research use only, not for use in diagnostic procedures.

DATA



Western blot analysis of PTEN phosphorylation in non-transfected: sc-117752 (A,D), untreated mouse PTEN transfected: sc-122834 (B,E) and lambda protein phosphatase (sc-200312A) treated mouse PTEN transfected: sc-122834 (C,F) 293T whole cell lysates. Antibodies tested include p-PTEN (H-3): sc-377573 (A,B,C) and PTEN (N-19): sc-6818 (D,E,F).



PTEN (N-19): sc-6818. Immunofluorescence staining of methanol-fixed HeLa cells showing nuclear localization.

SELECT PRODUCT CITATIONS

- Lu, Y., et al. 1999. The PTEN/MMAC1/TEP tumor suppressor gene decreases cell growth and induces apoptosis and anoikis in breast cancer cells. *Oncogene* 18: 7034-7045.
- Adachi, J., et al. 1999. Cell cycle arrest and astrocytic differentiation resulting from PTEN expression in glioma cells. *J. Neurosurg.* 91: 822-830.
- Kyrylenko, S., et al. 1999. Regulation of PTEN expression in neuronal apoptosis. *Brain Res. Mol. Brain Res.* 73: 198-202.
- Chunhu, Z., et al. 2008. Antiproliferative and apoptotic effects of paeonol on human hepatocellular carcinoma cells. *Anticancer Drugs* 19: 401-409.
- Mosessian, S., et al. 2009. Analysis of PTEN complex assembly and identification of heterogeneous nuclear ribonucleoprotein C as a component of the PTEN-associated complex. *J. Biol. Chem.* 284: 30159-30166.
- Cao, J., et al. 2009. Prdx1 inhibits tumorigenesis via regulating PTEN/AKT activity. *EMBO J.* 28: 1505-1517.
- Su, L., et al. 2010. EGR1 reactivation by histone deacetylase inhibitors promotes synovial sarcoma cell death through the PTEN tumor suppressor. *Oncogene* 29: 4352-4361.
- He, J., et al. 2010. The p85β regulatory subunit of PI3K serves as a substrate for PTEN protein phosphatase activity during Insulin mediated signaling. *Biochem. Biophys. Res. Commun.* 397: 513-519.
- Ochi, A., et al. 2012. Toll-like receptor 7 regulates pancreatic carcinogenesis in mice and humans. *J. Clin. Invest.* 122: 4118-4129.


 MONOS
Satisfaction
Guaranteed

Try **PTEN (A2B1): sc-7974** or **PTEN (F-1): sc-393186**, our highly recommended monoclonal alternatives to PTEN (N-19). Also, for AC, HRP, FITC, PE, Alexa Fluor® 488 and Alexa Fluor® 647 conjugates, see **PTEN (A2B1): sc-7974**.