

MAN2A1 (L-17): sc-68242

BACKGROUND

The α -mannosidases (designated MAN1A1, MAN1A2, MAN2A1 and MAN2A2) comprise a group of soluble proteins that localize to the endoplasmic reticulum, the Golgi apparatus or the cytoplasm. Depending on their cellular location, these proteins are involved in either the processing or the degradation of newly synthesized N-glycans. MAN2A1 (mannosidase α class 2A member 1) is a single-pass type II membrane protein that localizes to the cisternae of the Golgi and is involved in protein modification pathways. More specifically, MAN2A1 uses zinc as a cofactor to catalyze the first committed step in the formation of N-glycans, namely the hydrolysis of the terminal α -D-mannose residues in the oligosaccharide Man5(GlcNAc)3.

REFERENCES

- Moremen, K.W. and Robbins, P.W. 1991. Isolation, characterization, and expression of cDNAs encoding murine α -mannosidase II, a Golgi enzyme that controls conversion of high mannose to complex N-glycans. *J. Cell Biol.* 115: 1521-1534.
- Misago, M., et al. 1995. Molecular cloning and expression of cDNAs encoding human α -mannosidase II and a previously unrecognized α -mannosidase IIx isozyme. *Proc. Natl. Acad. Sci. USA* 92: 11766-11770.
- Chui, D., et al. 1997. α -mannosidase-II deficiency results in dyserythropoiesis and unveils an alternate pathway in oligosaccharide biosynthesis. *Cell* 90: 157-167.
- Chui, D., et al. 2001. Genetic remodeling of protein glycosylation *in vivo* induces autoimmune disease. *Proc. Natl. Acad. Sci. USA* 98: 1142-1147.
- Hart, M.L., et al. 2003. Glycosylation inhibitors and neuraminidase enhance human immunodeficiency virus type 1 binding and neutralization by mannose-binding lectin. *J. Gen. Virol.* 84: 353-360.
- Liu, T., et al. 2005. Human plasma N-glycoproteome analysis by immunoaffinity subtraction, hydrazide chemistry, and mass spectrometry. *J. Proteome Res.* 4: 2070-2080.

CHROMOSOMAL LOCATION

Genetic locus: MAN2A1 (human) mapping to 5q21.3; Man2a1 (mouse) mapping to 17 E1.1.

SOURCE

MAN2A1 (L-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of MAN2A1 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-68242 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

MAN2A1 (L-17) is recommended for detection of MAN2A1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

MAN2A1 (L-17) is also recommended for detection of MAN2A1 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for MAN2A1 siRNA (h): sc-61926, MAN2A1 siRNA (m): sc-61927, MAN2A1 shRNA Plasmid (h): sc-61926-SH, MAN2A1 shRNA Plasmid (m): sc-61927-SH, MAN2A1 shRNA (h) Lentiviral Particles: sc-61926-V and MAN2A1 shRNA (m) Lentiviral Particles: sc-61927-V.


Molecular Weight of MAN2A1: 131 kDa.

Positive Controls: mouse brain extract: sc-2253.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



MAN2A1 (L-17): sc-68242. Western blot analysis of MAN2A1 expression in mouse brain tissue extract.

RESEARCH USE

For research use only, not for use in diagnostic procedures.



Try **MAN2A1 (D-5): sc-377204** or **MAN2A1 (F-10): sc-376909**, our highly recommended monoclonal alternatives to MAN2A1 (L-17).