

Codanin-1 (H-301): sc-68404

BACKGROUND

The congenital dyserythropoietic anemias (CDAs) are an uncommon and heterogeneous group of disorders that are characterized by markedly ineffective erythropoiesis and, usually, striking dysplastic changes in erythroblasts. Congenital dyserythropoietic anemia type 1 (CDA1) is a rare autosomal recessive disorder with ineffective erythropoiesis, characteristic morphological abnormalities of erythroblasts and iron overloading. CDA1 is caused by mutations in the CDAN1 gene, which maps to chromosome 15q15.2 and encodes the 1,227 amino acid protein Codanin-1. Codanin-1 has a 150 residue N-terminal domain with sequence similarity to collagens and two shorter segments that show weak similarities to the microtubule-associated proteins synapsin and MAP-1B (neuraxin). Research indicates that Codanin-1 may be involved in nuclear envelope integrity, conceivably related to microtubule attachments. Skeletal anomalism has been associated with mutations of CDAN1, indicating that Codanin-1 may play a role in the development of the skeleton.

REFERENCES

1. Tamary, H., et al. 1996. Clinical features and studies of erythropoiesis in Israeli Bedouins with congenital dyserythropoietic anemia type I. *Blood* 87: 1763-1770.
2. Dgany, O., et al. 2002. Congenital dyserythropoietic anemia type 1 is caused by mutations in Codanin-1. *Am. J. Hum. Genet.* 71: 1467-1474.
3. Pielage, J., et al. 2003. The *Drosophila* cell survival gene discs lost encodes a cytoplasmic Codanin-1-like protein, not a homolog of tight junction PDZ protein Patj. *Dev. Cell* 5: 841-851.
4. Delaunay, J. 2003. Red cell membrane and erythropoiesis genetic defects. *Hematol. J.* 4: 225-232.
5. Heimpel, H., et al. 2005. Congenital dyserythropoietic anemia type 1 (CDA1): molecular genetics, clinical appearance, and prognosis based on long-term observation. *Blood* 107: 334-340.
6. Ahmed, M.R., et al. 2006. Evidence of genetic heterogeneity in congenital dyserythropoietic anaemia type I. *Br. J. Haematol.* 133: 444-446.
7. Ahmed, M.R., et al. 2006. Linkage and mutational analysis of the CDAN1 gene reveals genetic heterogeneity in congenital dyserythropoietic anemia type I. *Blood* 107: 4968-4969.
8. Goede, J.S., et al. 2006. Congenital dyserythropoietic anemia type 1 with bone abnormalities, mutations of the CDAN1 gene, and significant responsiveness to α -interferon therapy. *Ann. Hematol.* 85: 591-595.

CHROMOSOMAL LOCATION

Genetic locus: CDAN1 (human) mapping to 15q15.2; Cdan1 (mouse) mapping to 2 E5.

SOURCE

Codanin-1 (H-301) is a rabbit polyclonal antibody raised against amino acids 780-1080 mapping near the C-terminus of Codanin-1 of human origin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

Codanin-1 (H-301) is recommended for detection of Codanin-1 of human, mouse and, to a lesser extent, rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Codanin-1 (H-301) is also recommended for detection of Codanin-1 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for Codanin-1 siRNA (h): sc-62132, Codanin-1 siRNA (m): sc-62133, Codanin-1 shRNA Plasmid (h): sc-62132-SH, Codanin-1 shRNA Plasmid (m): sc-62133-SH, Codanin-1 shRNA (h) Lentiviral Particles: sc-62132-V and Codanin-1 shRNA (m) Lentiviral Particles: sc-62133-V.

Molecular Weight of Codanin-1: 134 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfation
Guaranteed

Try **Codanin-1 (G-1): sc-365839**, our highly recommended monoclonal alternative to Codanin-1 (H-301).