



ECA39 (S-17): sc-68659

BACKGROUND

Class-IV pyridoxal-phosphate-dependent aminotransferase family members ECA39 and BCAT2 are both enzymes that catalyze the first reaction in the catabolism of the essential branched chain amino acids valine, leucine and isoleucine. ECA39, also known as BCAT1 (branched-chain-amino-acid aminotransferase 1, cytosolic) is localized to the cytoplasm where it forms a homodimer. ECA39 is expressed in the brain and kidney during embryogenesis and is overexpressed in c-Myc induced tumors. BCAT2 (branched-chain-amino-acid aminotransferase 2, mitochondrial), also known as placental protein 18 (PP18), is expressed as two isoforms produced by alternative splicing. The first isoform of BCAT2, designated BCAT2A, is expressed in the mitochondrion, while the second isoform, designated BCAT2B, is expressed in the cytoplasm. Ubiquitously expressed, BCAT2 is also thought to act as a transporter of branched chain α -keto acids.

REFERENCES

- Schuldiner, O., Eden, A., Ben-Yosef, T., Yanuka, O., Simchen, G. and Benvenisty, N. 1996. ECA39, a conserved gene regulated by c-Myc in mice, is involved in G₁/S cell cycle regulation in yeast. *Proc. Natl. Acad. Sci. USA* 93: 7143-7148.
- Ben-Yosef, T., Eden, A. and Benvenisty, N. 1998. Characterization of murine BCAT genes: BCAT1, a c-Myc target, and its homolog, BCAT2. *Mamm. Genome*. 9: 595-597.
- Eden, A. and Benvenisty, N. 1999. Involvement of branched-chain amino acid aminotransferase (BCAT1/ ECA39) in apoptosis. *FEBS Lett.* 457: 255-261.
- Grimm, C.H., Rogner, U.C. and Avner, P. 2003. Lrmp and BCAT1 are candidates for the type I diabetes susceptibility locus Idd6. *Autoimmunity*. 36: 241-246.
- Yoshikawa, R., Yanagi, H., Shen, C.S., Fujiwara, Y., Noda, M., Yagyu, T., Gega, M., Oshima, T., Yamamura, T., Okamura, H., Nakano, Y., Morinaga, T. and Hashimoto-Tamaoki, T. 2006. ECA39 is a novel distant metastasis-related biomarker in colorectal cancer. *World J. Gastroenterol.* 12: 5884-5889.
- Zhou, W., Feng, X., Li, H., Wang, L., Li, H., Zhu, B., Zhang, H., Yao, K. and Ren, C. 2007. Functional evidence for a nasopharyngeal carcinoma-related gene BCAT1 located at 12p12. *Oncol. Res.* 16: 405-413.
- Conway, M.E., Coles, S.J., Islam, M.M. and Hutson, S.M. 2008. Regulatory control of human cytosolic branched-chain aminotransferase by oxidation and S-glutathionylation and its interactions with redox sensitive neuronal proteins. *Biochemistry* 47: 5465-5479.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

SOURCE

ECA39 (S-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of ECA39 of rat origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-68659 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

ECA39 (S-17) is recommended for detection of ECA39 of rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Molecular Weight of ECA39: 43 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RESEARCH USE

For research use only, not for use in diagnostic procedures.