

SAP 62 (N-17): sc-68793

BACKGROUND

SAP 62, also known as SF3A2 (splicing factor 3A subunit 2), PRP11, PRPF11 or SF3a66, is a 464 amino acid protein that contains one matrin-type zinc finger and belongs to the SF3A2 family. Localized to the nucleus, SAP 62 is a subunit of the SF3A splicing factor, a heterotrimeric complex comprised of three subunits that act in tandem to mediate the binding of U2 snRNP to the branchpoint sequence (BPS) in pre-mRNA. The SF3A complex is necessary for the conversion of 15S U2 snRNP into the active 17S protein that performs directly in pre-mRNA splicing events. Functioning as the second subunit of the complex, SAP 62 interacts with subunit 1 (SAP 114) via its N-terminus while simultaneously binding to 15S U2 snRNP via its zinc finger domain. In addition to its role in RNA splicing, SAP 62 is thought to act independently as a microtubule-binding protein.

REFERENCES

- Bennett, M. and Reed, R. 1993. Correspondence between a mammalian spliceosome component and an essential yeast splicing factor. *Science* 262: 105-108.
- Dresser, D.W., Hacker, A., Lovell-Badge, R. and Guerrier, D. 1995. The genes for a spliceosome protein (SAP62) and the anti-Müllerian hormone (AMH) are contiguous. *Hum. Mol. Genet.* 4: 1613-1618.
- Das, R., Zhou, Z. and Reed, R. 2000. Functional association of U2 snRNP with the ATP-independent spliceosomal complex E. *Mol. Cell* 5: 779-787.
- Dresser, D.W., Jamin, S.P., Atkins, C.J. and Guerrier, D. 2001. An expressed GNRP-like gene shares a bi-directional promoter with SF3A2 (SAP62) immediately upstream of AMH. *Gene* 277: 163-173.
- Jurica, M.S., Licklider, L.J., Gygi, S.R., Grigorieff, N. and Moore, M.J. 2002. Purification and characterization of native spliceosomes suitable for three-dimensional structural analysis. *RNA* 8: 426-439.
- Takenaka, K., Nakagawa, H., Miyamoto, S. and Miki, H. 2004. The pre-mRNA-splicing factor SF3a66 functions as a microtubule-binding and bundling protein. *Biochem. J.* 382: 223-230.
- Tanackovic, G. and Krämer, A. 2005. Human splicing factor SF3a, but not SF1, is essential for pre-mRNA splicing *in vivo*. *Mol. Biol. Cell* 16: 1366-1377.
- Oppelt, P., Strissel, P.L., Kellermann, A., Seeber, S., Humeny, A., Beckmann, M.W. and Strick, R. 2005. DNA sequence variations of the entire anti-Müllerian hormone (AMH) gene promoter and AMH protein expression in patients with the Mayer-Rokitanski-Kuster-Hauser syndrome. *Hum. Reprod.* 20: 149-157.
- Halm, S., Rocha, A., Miura, T., Prat, F. and Zanuy, S. 2007. Anti-Müllerian hormone (AMH/AMH) in the European sea bass: its gene structure, regulatory elements, and the expression of alternatively-spliced isoforms. *Gene* 388: 148-158.

CHROMOSOMAL LOCATION

Genetic locus: SF3A2 (human) mapping to 19p13.3; Sf3a2 (mouse) mapping to 10 C1.

SOURCE

SAP 62 (N-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of SAP 62 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-68793 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-68793 X, 200 µg/0.1 ml.

APPLICATIONS

SAP 62 (N-17) is recommended for detection of SAP 62 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

SAP 62 (N-17) is also recommended for detection of SAP 62 in additional species, including canine, bovine, porcine and avian.

Suitable for use as control antibody for SAP 62 siRNA (h): sc-76445, SAP 62 siRNA (m): sc-76446, SAP 62 shRNA Plasmid (h): sc-76445-SH, SAP 62 shRNA Plasmid (m): sc-76446-SH, SAP 62 shRNA (h) Lentiviral Particles: sc-76445-V and SAP 62 shRNA (m) Lentiviral Particles: sc-76446-V.

SAP 62 (N-17) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Molecular Weight of SAP 62: 66 kDa.

Positive Controls: HeLa nuclear extract: sc-2120 or A-431 nuclear extract: sc-2122.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.