

RASSF3 (H-60): sc-68854

BACKGROUND

Ras is a small GTP-binding protein involved in many cellular processes, including proliferation, differentiation and apoptosis. Ras transmits signals of cell surface receptors by binding to a variety of effector molecules. In addition to the well characterized effectors Raf and PI-3 kinase, Ras also interacts with members of the RASSF family, including RASSF1, RASSF2, RASSF3, RASSF4 and Nore1. Members of the RASSF family contain a highly conserved Ras association domain (RalGDS/AF-6 or RA) and function as Ras effectors/tumor suppressors. RASSF3 (Ras association domain family 3) is an ubiquitously expressed protein found in normal and cancerous tissues. RASSF3 contains an N-terminal RA domain and a coiled-coil SARAH domain. Three isoforms, namely RASSF3A, RASSF3B and RASSF3C, may exist for RASSF3 due to alternative splicing. RASSF3B and RASSF3C are shorter than RASSF3A and do not contain the RA and SARAH domain.

CHROMOSOMAL LOCATION

Genetic locus: RASSF3 (human) mapping to 12q14.2; RASSF3 (mouse) mapping to 10 D2.

SOURCE

RASSF3 (H-60) is a rabbit polyclonal antibody raised against amino acids 21-80 mapping near the N-terminus of RASSF3 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

RASSF3 (H-60) is recommended for detection of RASSF3 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

RASSF3 (H-60) is also recommended for detection of RASSF3 in additional species, including bovine.

Suitable for use as control antibody for RASSF3 siRNA (h): sc-62926, RASSF3 siRNA (m): sc-62927, RASSF3 shRNA Plasmid (h): sc-62926-SH, RASSF3 shRNA Plasmid (m): sc-62927-SH, RASSF3 shRNA (h) Lentiviral Particles: sc-62926-V and RASSF3 shRNA (m) Lentiviral Particles: sc-62927-V.


Molecular Weight RASSF3: 29 kDa.

Positive Controls: RASSF3 (m2): 293T Lysate: sc-122984.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



RASSF3 (H-60): sc-68854. Western blot analysis of RASSF3 expression in non-transfected: sc-117752 (A) and mouse RASSF3 transfected: sc-122984 (B) 293T whole cell lysates.



RASSF3 (H-60): sc-68854. Western blot analysis of RASSF3 expression in non-transfected: sc-117752 (A) and mouse RASSF3 transfected: sc-122983 (B) 293T whole cell lysates.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **RASSF3 (E-4): sc-376560**, our highly recommended monoclonal alternative to RASSF3 (H-60).