

NALP14 (H-95): sc-68858

BACKGROUND

NALP14 (NACHT, LRR and PYD domains-containing protein 14) is a testis specific nucleotide binding protein. It is a member of the NLRP family and contains one DAPIN (Pyrin, AIM, ASC and death domain-like (PAAD)) domain, eleven LRR (leucine-rich) repeats and one NACHT domain. NALP proteins most often act as an intracellular receptors involved in innate immunity. NALP14 has been linked to spermatogenesis but more likely functions as an inflammasome component. The inflammasome comprises a NALP member and an ASC adapter that ensures caspase-1 recruitment to the inflammasome. Different types of inflammasomes are stimulated by a wide variety of microbial, foreign and endogenous materials.

REFERENCES

1. Chamaillard, M., Girardin, S.E., Viala, J. and Philpott, D.J. 2003. Nods, Nalps and Naip: intracellular regulators of bacterial-induced inflammation. *Cell. Microbiol.* 5: 581-592.
2. Tschopp, J., Martinon, F. and Burns, K. 2003. NALPs: a novel protein family involved in inflammation. *Nat. Rev. Mol. Cell Biol.* 4: 95-104.
3. Horikawa, M., Kirkman, N.J., Mayo, K.E., Mulders, S.M., Zhou, J., Bondy, C.A., Hsu, S.Y., King, G.J. and Adashi, E.Y. 2005. The mouse germ-cell-specific leucine-rich repeat protein NALP1 a member of the NACHT nucleotide triphosphatase family. *Biol. Reprod.* 72: 879-889.
4. Kufer, T.A., Fritz, J.H. and Philpott, D.J. 2005. NACHT-LRR proteins (NLRs) in bacterial infection and immunity. *Trends Microbiol.* 13: 381-388.
5. Westerveld, G.H., Korver, C.M., van Pelt, A.M., Leschot, N.J., van der Veen, F., Repping, S. and Lombardi, M.P. 2006. Mutations in the testis-specific NALP14 gene in men suffering from spermatogenic failure. *Hum. Reprod.* 21: 3178-3184.
6. Martinon, F. 2007. Orchestration of pathogen recognition by inflammasome diversity: Variations on a common theme. *Eur. J. Immunol.* 37: 3003-3006.
7. Pétrilli, V. and Martinon, F. 2007. The inflammasome, autoinflammatory diseases, and gout. *Joint Bone Spine.* 74: 571-576.
8. Martinon, F., Gaide, O., Pétrilli, V., Mayor, A. and Tschopp, J. 2007. NALP inflammasomes: a central role in innate immunity. *Semin. Immunopathol.* 29: 213-229.

CHROMOSOMAL LOCATION

Genetic locus: NLRP14 (human) mapping to 11p15.4.

SOURCE

NALP14 (H-95) is a rabbit polyclonal antibody raised against amino acids 999-1093 mapping at the C-terminus of NALP14 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

NALP14 (H-95) is recommended for detection of NALP14 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for NALP14 siRNA (h): sc-62659, NALP14 shRNA Plasmid (h): sc-62659-SH and NALP14 shRNA (h) Lentiviral Particles: sc-62659-V.

Molecular Weight of NALP14: 125 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
 Satisfation
 Guaranteed

Try **NALP14 (H-10): sc-398068**, our highly recommended monoclonal alternative to NALP14 (H-95).