

ERR β (H-70): sc-68879

BACKGROUND

Estrogen and progesterone receptors are members of a family of transcription factors that are regulated by the binding of their cognate ligands. The interaction of hormone-bound estrogen receptors with estrogen responsive elements (EREs) alters transcription of ERE-containing genes. Estrogen receptor-related proteins (ERR α , β and γ) are orphan nuclear receptors. Like estrogen receptors, ERRs bind specifically to EREs to activate reporter genes. ERR β , also known as steroid hormone receptor ERR2 or estrogen receptor-like 2, contains a DNA binding domain and is highly homologous to ERR α . ERR β is expressed during mammary gland development and is critical in embryo development. It is expressed in a subset of diploid trophoblast cells which make up the chorion. The loss of ERR β results in severely impaired chorion formation leading to placental failure and embryonic death. This suggests that ERR β may be necessary for the proper formation or function of the chorion. In addition, ERR β potently represses the transcriptional activity of Nrf2.

REFERENCES

1. Luo, J., et al. 1997. Placental abnormalities in mouse embryos lacking the orphan nuclear receptor ERR β . *Nature* 388: 778-782.
2. Hong, H., et al. 1999. Hormone-independent transcriptional activation and coactivator binding by novel orphan nuclear receptor ERR3. *J. Biol. Chem.* 274: 22618-22626.
3. Chen, F., et al. 1999. Identification of two hERR2-related novel nuclear receptors utilizing bioinformatics and inverse PCR. *Gene* 228: 101-109.
4. Greschik, H., et al. 2002. Structural and functional evidence for ligand-independent transcriptional activation by the estrogen-related receptor 3. *Mol. Cell* 9: 303-313.
5. Cheung, C.P., et al. 2005. Expression and functional study of estrogen receptor-related receptors in human prostatic cells and tissues. *J. Clin. Endocrinol. Metab.* 90: 1830-1844.
6. Gearhart, M.D., et al. 2005. Inhibition of DNA binding by human estrogen-related receptor 2 and estrogen receptor α with minor groove binding polyamides. *Biochemistry* 44: 4196-4203.

CHROMOSOMAL LOCATION

Genetic locus: ESRRB (human) mapping to 14q24.3.

SOURCE

ERR β (H-70) is a rabbit polyclonal antibody raised against amino acids 431-500 mapping at the C-terminus of ERR β of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4 $^{\circ}$ C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

ERR β (H-70) is recommended for detection of ERR β of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for ERR β siRNA (h): sc-60603, ERR β shRNA Plasmid (h): sc-60603-SH and ERR β shRNA (h) Lentiviral Particles: sc-60603-V.


Molecular Weight of ERR β : 56 kDa.

Positive Controls: ERR β (h): 293T Lysate: sc-128549 or A549 cell lysate: sc-2413.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

DATA



ERR β (H-70): sc-68879. Western blot analysis of ERR β expression in non-transfected 293T: sc-117752 (A), human ERR β transfected 293T: sc-128549 (B) and A549 (C) whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **ERR β / γ (E-1): sc-376449**, our highly recommended monoclonal alternative to ERR β (H-70).