

ST8Sia IV (H-70): sc-68896

BACKGROUND

ST8Sia IV (ST8 α -N-acetyl-neuraminide α -2,8-sialyltransferase IV), also known as PST, PST1 or SIAT8D, is a 359 amino acid single-pass type II membrane protein that localizes to the membrane of the Golgi apparatus. Highly expressed in heart, thymus and spleen, as well as fetal lung, brain and kidney, ST8Sia IV functions to catalyze the polycondensation of α -2,8-linked sialic acid, an event that is required for the synthesis of polysialic acid (PSA). PSA is an important regulator of neuronal plasticity and is present in embryonic brain tissue, where it interacts with NCAM (neural cell adhesion molecule) and plays a crucial role in fetal brain development. Defects in the gene encoding ST8Sia IV are associated with idiopathic pancreatitis, schizophrenia and tumor formation/metastasis. ST8Sia IV exists as multiple isoforms produced by alternative splicing events.

REFERENCES

1. Nakayama, J., Fukuda, M.N., Fredette, B., Ranscht, B. and Fukuda, M. 1995. Expression cloning of a human polysialyltransferase that forms the polysialylated neural cell adhesion molecule present in embryonic brain. *Proc. Natl. Acad. Sci. USA* 92: 7031-7035.
2. Angata, K., Yen, T.Y., El-Battari, A., Macher, B.A. and Fukuda, M. 2001. Unique disulfide bond structures found in ST8Sia IV polysialyltransferase are required for its activity. *J. Biol. Chem.* 276: 15369-15377.
3. Angata, K., Suzuki, M. and Fukuda, M. 2002. ST8Sia II and ST8Sia IV polysialyltransferases exhibit marked differences in utilizing various acceptors containing oligosialic acid and short polysialic acid. The basis for cooperative polysialylation by two enzymes. *J. Biol. Chem.* 277: 36808-36817.
4. Cohn, J.A., Noone, P.G. and Jowell, P.S. 2002. Idiopathic pancreatitis related to CFTR: complex inheritance and identification of a modifier gene. *J. Investig. Med.* 50: 247S-255S.
5. Beecken, W.D., Engl, T., Ogbomo, H., Relja, B., Cinatl, J., Bereiter-Hahn, J., Oppermann, E., Jonas, D. and Blaheta, R.A. 2005. Valproic acid modulates NCAM polysialylation and polysialyltransferase mRNA expression in human tumor cells. *Int. Immunopharmacol.* 5: 757-769.
6. Mendiratta, S.S., Sekulic, N., Lavie, A. and Colley, K.J. 2005. Specific amino acids in the first fibronectin type III repeat of the neural cell adhesion molecule play a role in its recognition and polysialylation by the polysialyltransferase ST8Sia IV/PST. *J. Biol. Chem.* 280: 32340-32348.

CHROMOSOMAL LOCATION

Genetic locus: ST8SIA4 (human) mapping to 5q21.1; St8sia4 (mouse) mapping to 1 D.

SOURCE

ST8Sia IV (H-70) is a rabbit polyclonal antibody raised against amino acids 21-90 mapping near the N-terminus of ST8Sia IV of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

ST8Sia IV (H-70) is recommended for detection of ST8Sia IV of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

ST8Sia IV (H-70) is also recommended for detection of ST8Sia IV in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for ST8Sia IV siRNA (h): sc-92058, ST8Sia IV siRNA (m): sc-153868, ST8Sia IV shRNA Plasmid (h): sc-92058-SH, ST8Sia IV shRNA Plasmid (m): sc-153868-SH, ST8Sia IV shRNA (h) Lentiviral Particles: sc-92058-V and ST8Sia IV shRNA (m) Lentiviral Particles: sc-153868-V.


Molecular Weight of ST8Sia IV: 46 kDa.

Positive Controls: mouse spleen extract : sc-2391, mouse brain extract: sc-2253 or ST8Sia IV (h): 293 Lysate: sc-114067.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



ST8Sia IV (H-70): sc-68896. Western blot analysis of ST8Sia IV expression in non-transfected: sc-110760 (A) and human ST8Sia IV transfected: sc-114067 (B) 293 whole cell lysates and mouse brain tissue extract (C).



ST8Sia IV (H-70): sc-68896. Western blot analysis of ST8Sia IV expression in non-transfected: sc-117752 (A) and human ST8Sia IV transfected: sc-116438 (B) 293T whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.