

V-ATPase D1 (N-16): sc-69106

BACKGROUND

Vacuolar-type H⁺-ATPase (V-ATPase) is a multisubunit enzyme responsible for the acidification of eukaryotic intracellular organelles. V-ATPases pump protons against an electrochemical gradient, while F-ATPases reverse the process, thereby synthesizing ATP. A peripheral V1 domain, which is responsible for ATP hydrolysis, and an integral V0 domain, which is responsible for proton translocation, comprise the V-ATPase complex. Nine subunits (A-H) make up the V1 domain and five subunits (A, D, C, C' and C'') make up the V0 domain. V-ATPase D1 (ATPase, H⁺ transporting, lysosomal, V0 subunit D1), also known as ATP6V0D1, P39, VATX, VMA6, ATP6D or VPATPD, is the D subunit of the V0 domain. Expressed ubiquitously, V-ATPase D1 acts in concert with other V0 subunits to catalytically acidify a variety of intracellular compartments, thereby synthesizing ATP to be used for vacuolar transport.

REFERENCES

1. van Hille, B., Vanek, M., Richener, H., Green, J.R. and Bilbe, G. 1993. Cloning and tissue distribution of subunits C, D, and E of the human vacuolar H⁺-ATPase. *Biochem. Biophys. Res. Commun.* 197: 15-21.
2. Finbow, M.E. and Harrison, M.A. 1997. The vacuolar H⁺-ATPase: a universal proton pump of eukaryotes. *Biochem. J.* 324: 697-712.
3. Forgac, M. 1999. Structure and properties of the vacuolar H⁺-ATPases. *J. Biol. Chem.* 274: 12951-12954.
4. Agarwal, A.K. and White, P.C. 2000. Structure of the VPATPD gene encoding subunit D of the human vacuolar proton ATPase. *Biochem. Biophys. Res. Commun.* 279: 543-547.
5. Smith, A.N., Borthwick, K.J. and Karet, F.E. 2002. Molecular cloning and characterization of novel tissue-specific isoforms of the human vacuolar H⁺-ATPase C, G and d subunits, and their evaluation in autosomal recessive distal renal tubular acidosis. *Gene* 297: 169-177.
6. Nishi, T. and Forgac, M. 2002. The vacuolar H⁺-ATPases—nature's most versatile proton pumps. *Nat. Rev. Mol. Cell Biol.* 3: 94-103.
7. Morel, N. 2003. Neurotransmitter release: the dark side of the vacuolar H⁺-ATPase. *Biol. Cell* 95: 453-457.
8. Kawasaki-Nishi, S., Nishi, T. and Forgac, M. 2003. Proton translocation driven by ATP hydrolysis in V-ATPases. *FEBS Lett.* 545: 76-85.
9. Pietrement, C., Sun-Wada, G.H., Silva, N.D., McKee, M., Marshansky, V., Brown, D., Futai, M. and Breton, S. 2006. Distinct expression patterns of different subunit isoforms of the V-ATPase in the rat epididymis. *Biol. Reprod.* 74: 185-194.

CHROMOSOMAL LOCATION

Genetic locus: ATP6V0D1 (human) mapping to 16q22.1; Atp6v0d1 (mouse) mapping to 8 D3.

SOURCE

V-ATPase D1 (N-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of V-ATPase D1 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-69106 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

V-ATPase D1 (N-16) is recommended for detection of V-ATPase D1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

V-ATPase D1 (N-16) is also recommended for detection of V-ATPase D1 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for V-ATPase D1 siRNA (h): sc-63207, V-ATPase D1 siRNA (m): sc-63208, V-ATPase D1 shRNA Plasmid (h): sc-63207-SH, V-ATPase D1 shRNA Plasmid (m): sc-63208-SH, V-ATPase D1 shRNA (h) Lentiviral Particles: sc-63207-V and V-ATPase D1 shRNA (m) Lentiviral Particles: sc-63208-V.

Molecular Weight of V-ATPase D1: 40 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **V-ATPase D1 (D-4): sc-393322** or **V-ATPase D1 (34-Z): sc-81887**, our highly recommended monoclonal alternatives to V-ATPase D1 (N-16).