

Nanos2 (A-12): sc-69187

BACKGROUND

Nanos2, also known as NOS2, is a 138 amino acid protein that contains one nanos-type zinc finger. The nanos-type zinc finger is comprised of 2 C₂HC motifs, each of which are capable of binding one molecule of zinc and each of which work in tandem to play essential roles in translational regulation events. Expressed specifically in male germ cells, Nanos2 is essential for spermatogonia formation and is required to support the self-renewal, proliferation and overall development of proximal germ cells. Additionally, Nanos2 is thought to regulate the translation of target mRNAs, possibly by associating with the 3'-UTR of select transcripts. The gene encoding human Nanos2 maps to chromosome 19, which is the genetic home for a number of immunoglobulin superfamily members, including the killer cell and leukocyte Ig-like receptors, a number of ICAMs, the CEACAM and PSG family and Fc receptors (FcRs).

REFERENCES

1. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 608228. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
2. Jaruzelska, J., et al. 2003. Conservation of a Pumilio-Nanos complex from *Drosophila* germ plasm to human germ cells. *Dev. Genes Evol.* 213: 120-126.
3. Tsuda, M., et al. 2003. Conserved role of Nanos proteins in germ cell development. *Science* 301: 1239-1241.
4. D'Agostino, I., et al. 2006. Translational repression restricts expression of the *C. elegans* Nanos homolog NOS-2 to the embryonic germline. *Dev. Biol.* 292: 244-252.
5. Tsuda, M., et al. 2006. Implication of Nanos2-3'UTR in the expression and function of Nanos2. *Mech. Dev.* 123: 440-449.
6. Suzuki, A., et al. 2007. Functional redundancy among Nanos proteins and a distinct role of Nanos2 during male germ cell development. *Development* 134: 77-83.
7. Saga, Y. 2008. Mouse germ cell development during embryogenesis. *Curr. Opin. Genet. Dev.* 18: 337-341.

CHROMOSOMAL LOCATION

Genetic locus: NANOS2 (human) mapping to 19q13.32; Nanos2 (mouse) mapping to 7 A2.

SOURCE

Nanos2 (A-12) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of Nanos2 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-69187 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

Nanos2 (A-12) is recommended for detection of Nanos2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Nanos2 (A-12) is also recommended for detection of Nanos2 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for Nanos2 siRNA (h): sc-75866, Nanos2 siRNA (m): sc-75867, Nanos2 shRNA Plasmid (h): sc-75866-SH, Nanos2 shRNA Plasmid (m): sc-75867-SH, Nanos2 shRNA (h) Lentiviral Particles: sc-75866-V and Nanos2 shRNA (m) Lentiviral Particles: sc-75867-V.

Molecular Weight of Nanos2: 18 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
 Satisfaction
 Guaranteed

Try **Nanos2 (B-12): sc-393794** or **Nanos2 (A-8): sc-393868**, our highly recommended monoclonal alternatives to Nanos2 (A-12).