COX4 (K-20): sc-69361



The Power to Question

BACKGROUND

Cytochrome c oxidase (COX) functions as the terminal oxidase of the respiratory chain that uses cytochrome c as an electron donor to drive a proton gradient across the inner mitochondrial membrane. The mammalian COX apoenzyme is a heteromer consisting of three mitochondrial encoded catalytic subunits and several nuclear gene encoded structural subunits. COX contains two iron-coordination sites and two copper-coordination sites. Cytochrome c oxidase IV (COX4) is a nuclear-encoded subunit of COX that may play a role in regulating COX activity. COX4 is expressed ubiquitously in adult human tissue with the strongest levels of expression in the pancreas and moderate expression levels in heart, skeletal muscle and placenta.

REFERENCES

- Steffens, G.J., et al. 1979. Studies on cytochrome c oxidase, IV[1-3]. Primary structure and function of subunit II. Hoppe-Seyler's Z. Physiol. Chem. 360: 613-619.
- Brown, W.M., et al. 1982. Mitochondrial DNA sequences of primates: tempo and mode of evolution. J. Mol. Evol. 18: 225-239.
- Zeviani, M., et al. 1987. Isolation of a cDNA clone encoding subunit IV of human cytochrome c oxidase. Gene 55: 205-217.
- Lomax, M.I., et al. 1992. Rapid evolution of the human gene for cytochrome c oxidase subunit IV. Proc. Natl. Acad. Sci. USA 89: 5266-5270.
- Makris, G.J., et al. 1997. The gene encoding subunit IV of cytochrome c oxidase maps to mouse chromosome 8. Mamm. Genome 7: 789-790.
- Bachman, N.J., et al. 1999. The 5' region of the COX4 gene contains a novel overlapping gene, NOC4. Mamm. Genome 10: 506-512.
- 7. Hüttemann, M., et al. 2001. Mammalian subunit IV isoforms of cytochrome c oxidase. Gene 267: 111-123.
- 8. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 123864. World Wide Web URL: http://www.ncbi.nlm.nih.gov/omim/
- 9. Vizirianakis, I.S., et al. 2002. Differentiation-dependent repression of c-myc, B22, COX II and COX IV genes in murine erythroleukemia (MEL) cells. Biochem. pharmacol. 63: 1009-1017.

CHROMOSOMAL LOCATION

Genetic locus: Cox4i1 (mouse) mapping to 8 E1.

SOURCE

COX4 (K-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of COX4 of mouse origin.

PRODUCT

Each vial contains 200 μg lgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-69361 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

COX4 (K-20) is recommended for detection of cytochrome c oxidase IV of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for COX4 siRNA (m): sc-72075, COX4 shRNA Plasmid (m): sc-72075-SH and COX4 shRNA (m) Lentiviral Particles: sc-72075-V.


Molecular Weight of COX4: 17 kDa.

Positive Controls: mouse brain extract: sc-2253 or mouse heart extract: sc-2254.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



COX4 (K-20): sc-69361. Western blot analysis of COX4 expression in mouse heart tissue extract.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3801 Fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com