

CREB3L3 (N-14): sc-69375

BACKGROUND

CREB3L3 (cAMP-responsive element-binding protein 3-like protein 3), also known as CREBH or HYST1481, is a 461 amino acid single-pass type II membrane protein that localizes to the endoplasmic reticulum (ER) and, in response to ER stress, is cleaved and translocated to the nucleus. Expressed exclusively in liver, CREB3L3 functions as a transcription factor that, during ER stress, is thought to activate genes that are involved in both the unfolded protein response and the acute phase response (APR). Additionally, CREB3L3 is under-expressed in hepatocellular carcinoma, suggesting a possible role as a tumor suppressor. CREB3L3 functions as a dimer and contains one leucine zipper domain, a KDEL-like sequence and a bZIP domain, through which it conveys its DNA binding ability. Three isoforms of CREB3L3 exist due to alternative splicing events.

CHROMOSOMAL LOCATION

Genetic locus: CREB3L3 (human) mapping to 19p13.3; Creb3l3 (mouse) mapping to 10 C1.

SOURCE

CREB3L3 (N-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of CREB3L3 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-69375 X, 200 µg/0.1 ml.

Blocking peptide available for competition studies, sc-69375 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

CREB3L3 (N-14) is recommended for detection of CREB3L3 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

CREB3L3 (N-14) is also recommended for detection of CREB3L3 in additional species, including canine, bovine and porcine.

Suitable for use as control antibody for CREB3L3 siRNA (h): sc-72999, CREB3L3 siRNA (m): sc-77022, CREB3L3 shRNA Plasmid (h): sc-72999-SH, CREB3L3 shRNA Plasmid (m): sc-77022-SH, CREB3L3 shRNA (h) Lentiviral Particles: sc-72999-V and CREB3L3 shRNA (m) Lentiviral Particles: sc-77022-V.


CREB3L3 (N-14) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Molecular Weight of CREB3L3: 50 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 3) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

DATA



CREB3L3 (N-14): sc-69375. Immunoperoxidase staining of formalin fixed, paraffin-embedded human duodenum tissue showing nuclear and cytoplasmic staining of glandular cells.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **CREB3L3 (E-2): sc-377332** or **CREB3L3 (G-11): sc-377156**, our highly recommended monoclonal alternatives to CREB3L3 (N-14).