

JMJD5 (Y-20): sc-69429

BACKGROUND

JMJD5 (Jumonji domain containing 5) is a nuclear protein that is believed to function as a histone lysine demethylase. Belonging to the Jumonji C-domain-containing histone lysine demethylase (JHDM) family, JMJD5 contains one JMJC (Jumonji C) domain. The *C. elegans* homolog of JMJD5 has been identified as a protein that protects the genome against insertions and deletions. This suggests a potential role for mammalian JMJD5 as a tumor suppressor. Further supporting the role of JMJD5 as a tumor suppressor, the knockdown of JMJD5 expression in mouse fibroblasts can lead to an increased mutation rate and an increased tolerance to MNNG (a DNA methylating agent). This implies that JMJD5 may specifically participate in DNA mismatch repair.

REFERENCES

1. Toyoda, M., Kojima, M. and Takeuchi, T. 2000. Jumonji is a nuclear protein that participates in the negative regulation of cell growth. *Biochem. Biophys. Res. Commun.* 274: 332-336.
2. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 611917. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
3. Jung, J., Mysliwiec, M.R. and Lee, Y. 2005. Roles of Jumonji in mouse embryonic development. *Dev. Dyn.* 232: 21-32.
4. Takeuchi, T., Watanabe, Y., Takano-Shimizu, T. and Kondo, S. 2006. Roles of Jumonji and Jumonji family genes in chromatin regulation and development. *Dev. Dyn.* 235: 2449-2459.
5. Suzuki, T., Minehata, K., Akagi, K., Jenkins, N.A. and Copeland, N.G. 2006. Tumor suppressor gene identification using retroviral insertional mutagenesis in Blm-deficient mice. *EMBO J.* 25: 3422-3431.
6. Cui, L., Fan, Q., Cui, L. and Miao, J. 2008. Histone lysine methyltransferases and demethylases in *Plasmodium falciparum*. *Int. J. Parasitol.* 38: 1083-1097.

CHROMOSOMAL LOCATION

Genetic locus: JMJD5 (human) mapping to 16p12.1; Jmjd5 (mouse) mapping to 7 F3.

SOURCE

JMJD5 (Y-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of JMJD5 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-69429 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

JMJD5 (Y-20) is recommended for detection of JMJD5 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

JMJD5 (Y-20) is also recommended for detection of JMJD5 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for JMJD5 siRNA (h): sc-75359, JMJD5 siRNA (m): sc-75360, JMJD5 shRNA Plasmid (h): sc-75359-SH, JMJD5 shRNA Plasmid (m): sc-75360-SH, JMJD5 shRNA (h) Lentiviral Particles: sc-75359-V and JMJD5 shRNA (m) Lentiviral Particles: sc-75360-V.


Molecular Weight of JMJD5: 47 kDa.

Positive Controls: JMJD5 (m): 293T Lysate: sc-127027.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



JMJD5 (Y-20): sc-69429. Western blot analysis of JMJD5 expression in non-transfected: sc-117752 (A) and mouse JMJD5 transfected: sc-127027 (B) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **JMJD5 (D-5): sc-377078** or **JMJD5 (G-3): sc-377440**, our highly recommended monoclonal alternatives to JMJD5 (Y-20).