

# HERP (S-18): sc-69497

## BACKGROUND

The endoplasmic reticulum (ER) stress response is triggered by the accumulation of unfolded proteins within the ER and is characterized by three events: the inhibition of translation (to prevent further protein accumulation), the up-regulated expression of polypeptide-folding proteins (known as the unfolded protein response, or UPR) and the degradation of misfolded proteins by the ER-associated protein degradation (ERAD) system. HERP, also known as HERPUD1 (homocysteine-inducible, endoplasmic reticulum stress-inducible, ubiquitin-like domain member 1), SUP or MIF1, is a 391 amino acid multi-pass membrane protein that localizes to the ER and contains one N-terminal ubiquitin-like domain. Widely expressed with highest expression in the brain, HERP is a component of the ERAD system and, via its ubiquitin-like domain, is thought to be involved in the destruction of misfolded proteins. Three isoforms of HERP exist due to alternative splicing events.

## CHROMOSOMAL LOCATION

Genetic locus: HERPUD1 (human) mapping to 16q13; Herpud1 (mouse) mapping to 8 C5.

## SOURCE

HERP (S-18) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of HERP of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-69497 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

HERP (S-18) is recommended for detection of HERP of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

HERP (S-18) is also recommended for detection of HERP in additional species, including canine, bovine and porcine.

Suitable for use as control antibody for HERP siRNA (h): sc-75245, HERP siRNA (m): sc-75246, HERP shRNA Plasmid (h): sc-75245-SH, HERP shRNA Plasmid (m): sc-75246-SH, HERP shRNA (h) Lentiviral Particles: sc-75245-V and HERP shRNA (m) Lentiviral Particles: sc-75246-V.

Molecular Weight of HERP: 54 kDa.

Positive Controls: Hep G2 cell lysate: sc-2227.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 3) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

## DATA



HERP (S-18): sc-69497. Immunoperoxidase staining of formalin fixed, paraffin-embedded human skeletal muscle tissue showing cytoplasmic staining of myocytes.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.

**MONOS**  
Satisfaction  
Guaranteed

Try **HERP (19-Y): sc-100721**, our highly recommended monoclonal alternative to HERP (S-18).