

# OS4 (C-19): sc-69518

## BACKGROUND

OS4, also known as CTDSP2 (carboxy-terminal domain RNA polymerase II polypeptide A small phosphatase 2) or SCP2, is a 271 amino acid nuclear protein that is implicated in the development of sarcomas. Expressed ubiquitously with particularly high expression observed in the pancreas, OS4 functions in the regulation of androgen-dependent transcription by globally silencing neuronal genes. OS4 contains one FCP1 homology domain and, acting as a Smad2/3 linker phosphatase, can enhance Smad2/3 signaling. Through its ability to interact with Smad2/3, OS4 contributes to the final signaling events in the Smad pathway, and thus is thought to be a crucial component in the metastasis of certain cancers. Overexpression of OS4 may lead to the development of primary sarcomas, as well as brain tumors.

## REFERENCES

1. Su, Y.A., Lee, M.M., Hutter, C.M. and Meltzer, P.S. 1997. Characterization of a highly conserved gene (OS4) amplified with Cdk4 in human sarcomas. *Oncogene* 15: 1289-1294.
2. Zohn, I.E. and Brivanlou, A.H. 2002. Expression cloning of *Xenopus* Os4, an evolutionarily conserved gene, which induces mesoderm and dorsal axis. *Dev. Biol.* 239: 118-131.
3. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 608711. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. Thompson, J., Lepikhova, T., Teixeira-Travesa, N., Whitehead, M.A., Palvimo, J.J. and Jänne, O.A. 2006. Small carboxyl-terminal domain phosphatase 2 attenuates androgen-dependent transcription. *EMBO J.* 25: 2757-2767.
5. Knockaert, M., Sapkota, G., Alarcón, C., Massagué, J. and Brivanlou, A.H. 2006. Unique players in the BMP pathway: small C-terminal domain phosphatases dephosphorylate Smad1 to attenuate BMP signaling. *Proc. Natl. Acad. Sci. USA* 103: 11940-11945.
6. Wrighton, K.H., Willis, D., Long, J., Liu, F., Lin, X. and Feng, X.H. 2006. Small C-terminal domain phosphatases dephosphorylate the regulatory linker regions of Smad2 and Smad3 to enhance transforming growth factor- $\beta$  signaling. *J. Biol. Chem.* 281: 38365-38375.
7. Yeo, M. and Lin, P.S. 2007. Functional characterization of small CTD phosphatases. *Methods Mol. Biol.* 365: 335-346.

## CHROMOSOMAL LOCATION

Genetic locus: CTDSP2 (human) mapping to 12q14.1; Ctdsp2 (mouse) mapping to 10 D3.

## SOURCE

OS4 (C-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of OS4 of human origin.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-69518 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

OS4 (C-19) is recommended for detection of OS4 of mouse and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

OS4 (C-19) is also recommended for detection of OS4 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for OS4 siRNA (h): sc-76008, OS4 siRNA (m): sc-155922, OS4 shRNA Plasmid (h): sc-76008-SH, OS4 shRNA Plasmid (m): sc-155922-SH, OS4 shRNA (h) Lentiviral Particles: sc-76008-V and OS4 shRNA (m) Lentiviral Particles: sc-155922-V.

Molecular Weight of OS4: 32 kDa.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.



Try **OS4 (2230C1a): sc-81347**, our highly recommended monoclonal alternative to OS4 (C-19).