SANTA CRUZ BIOTECHNOLOGY, INC.

Mnk1 (M-20): sc-6962



BACKGROUND

The MAPKAP kinases (for MAP kinase activated protein kinases) are a group of MAP kinase substrates which are themselves kinases. In response to activation, the MAP kinases phosphorylate downstream components on a consensus Pro-X-Ser/Thr-Pro motif. Several kinases that contain this motif have been identifed and serve as substrates for the ERK and p38 MAP kinases. These include the serine/threonine kinases Rsk-1 (also designated MAPKAP kinase-1), Rsk-2 and Rsk-3, which are phosphorylated by ERK 1 and ERK 2. Similarly p38 phosphorylates and activates the serine/threonine kinases MAPKAP kinase-2 and MAPKAP kinase-3 (also designated 3pK). The serine/threonine kinases Mnk1 and Mnk2 are substrates for both ERK and p38 MAP kinases.

REFERENCES

- 1. Sturgill, T.W., et al. 1988. Insulin-stimulated MAP2 kinase phosphorylates and activates ribosomal protein S6 kinase II. Nature 334: 715-718.
- Stokoe, D., et al. 1992. MAPKAP kinase-2: a novel protein kinase activated by mitogen-activated protein kinase. EMBO J. 11: 3985-3994.
- Davis, R.J. 1993. The mitogen-activated protein kinase signal transduction pathway. J. Biol. Chem. 268: 14553-14556.
- Zhao, Y., et al. 1995. RSK3 encodes a novel pp90Rsk isoform with a unique N-terminal sequence: growth factor stimulated kinase function and nuclear translocation. Mol. Cell. Biol. 15: 4353-4363.
- McLaughlin, M.M., et al. 1996. Identification of mitogen-activated protein (MAP) kinase-activated protein kinase-3, a novel substrate of CSBP p38 MAP kinase. J. Biol. Chem. 271: 8488-8492.
- Sithanandam, G., et al. 1996. 3pK, a new mitogen-activated protein kinaseactivated protein kinase located in the small cell lung cancer tumor suppressor gene region. Mol. Cell. Biol. 16: 868-876.
- Fukunaga, R. and Hunter, T. 1997. Mnk1, a new MAP kinase-activated protein kinase, isolated by a novel expression screening method for identifying protein kinase substrates. EMBO J. 16: 1921-1933.

CHROMOSOMAL LOCATION

Genetic locus: MKNK1 (human) mapping to 1p33; Mknk1 (mouse) mapping to 4 D1.

SOURCE

Mnk1 (M-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of Mnk1 of mouse origin.

PRODUCT

Each vial contains 200 μg lgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-6962 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS


Mnk1 (M-20) is recommended for detection of Mnk1 of mouse, rat and, to a lesser extent, human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Mnk1 siRNA (h): sc-39106, Mnk1 siRNA (m): sc-39107, Mnk1 shRNA Plasmid (h): sc-39106-SH, Mnk1 shRNA Plasmid (m): sc-39107-SH, Mnk1 shRNA (h) Lentiviral Particles: sc-39106-V and Mnk1 shRNA (m) Lentiviral Particles: sc-39107-V.

Molecular Weight of Mnk1: 52 kDa.

Positive Controls: Mnk1 (m): 293T Lysate: sc-125627 or 3611-RF whole cell lysate: sc-2215.

DATA



Mnk1 (M-20): sc-6962. Western blot analysis of Mnk1 expression in non-transfected: sc-11752 (**A**) and mouse Mnk1 transfected: sc-125627 (**B**) 293T whole cell lysates.

SELECT PRODUCT CITATIONS

- 1. Hsieh, C.C., et al. 2004. Akt/PKB and p38 MAPK signaling, translational initiation and longevity in Snell dwarf mouse livers. Mech. Ageing Dev. 125: 785-798.
- 2. DeWire, S.M., et al. 2008. β -Arrestin-mediated signaling regulates protein synthesis. J. Biol. Chem. 283: 10611-10620.

STORAGE

Store at 4° C, **D0 NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS Satisfation

Guaranteed

Try Mnk1 (A-4): sc-133107 or Mnk1 (C-5): sc-133108, our highly recommended monoclonal alternatives to Mnk1 (M-20).