



Nectin 3 (N3.2.9): sc-69714

BACKGROUND

Nectin is a Ca^{2+} -independent homophilic cell adhesion molecule that belongs to the immunoglobulin superfamily. Human Nectin is identical to the poliovirus receptor-related protein (PRR) and is identified to be the α herpesvirus entry mediator. Nectin constitutes a family consisting of at least Nectin 1, 2 and 3. Nectin 2 and 3 are ubiquitously expressed, whereas Nectin 1 is abundantly expressed in brain. Nectin 3, also designated as PRR3, has three splicing variants: Nectin 3 α , 3 β and 3 γ . Nectin 3 is a transmembrane protein whose extracellular region contains three Ig-like domains. Nectin 3 α and 3 β , but not Nectin 3 γ , have a C-terminal conserved motif (E/A-X-Y-V). This motif interacts with the PDZ domain of the F-Actin-binding protein, afadin, through which it is linked to the Actin cytoskeleton. Nectin 3 α co-localizes with Nectin 2 at adherin-based adheren junctions and interacts with afadin.

REFERENCES

- Lopez, M., Eberie, F., Mattei, M.G., Gabert, J., Birg, F., Bardin, F., Maroc, C. and Dubreuil, P. 1995. Complementary DNA characterization and chromosomal localization of a human gene related to the poliovirus receptor-encoding gene. *Gene* 155: 261-265.
- Asakura, T., Nakanishi, H., Sakisaka, T., Takahashi, K., Mandai, K., Nishimura, M., Sasaki, T. and Takai, T. 1999. Similar and differential behavior between the nectin-afadin-ponsin and cadherin-catenin systems during the formation and disruption of the polarized junctional alignment in epithelial cells. *Genes Cells* 4: 573-581.
- Kikyo, M., Matozaki, T., Kodama, A., Kawabe, H., Nakanishi, H. and Takai, Y. 2000. Cell-cell adhesion-mediated tyrosine phosphorylation of Nectin 2 δ , an immunoglobulin-like cell adhesion molecule at adherens junctions. *Oncogene* 19: 4022-4028.
- Satoh-Horikawa, K., Nakanishi, H., Takahashi, K., Miyahara, M., Nishimura, M., Tachibana, K., Mizoguchi, A. and Takai, Y. 2000. Nectin 3, a new member of immunoglobulin-link cell adhesion molecules that shows homophilic and heterophilic cell-cell adhesion activities. *J. Biol. Chem.* 275: 10291-10299.
- Reymond, N., Borg, J.P., Lecocq, E., Adalaide, J., Campadelli-Fiume, G., Dubreuil, P. and Lopez, M. 2000. Human Nectin 3/PRR3: a novel member of the PVR/PRR/Nectin family that interacts with afadin. *Gene* 255: 347-355.

CHROMOSOMAL LOCATION

Genetic locus: PVRL3 (human) mapping to 3q13.

SOURCE

Nectin 3 (N3.2.9) is a mouse monoclonal antibody raised against full-length recombinant Nectin 3 of human origin.

PRODUCT

Each vial contains 200 μg IgG_{2a} in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Nectin 3 (N3.2.9) is available conjugated to either phycoerythrin (sc-69714 PE) or fluorescein (sc-69714 FITC), 200 $\mu\text{g}/\text{ml}$, for WB (RGB), IF, IHC(P) and FCM.

APPLICATIONS

Nectin 3 (N3.2.9) is recommended for detection of Nectin 3 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μg per 100-500 μg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and flow cytometry (1 μg per 1×10^6 cells); may cross-react with Nectin 3 of monkey origin.

Suitable for use as control antibody for Nectin 3 siRNA (h): sc-36024, Nectin 3 shRNA Plasmid (h): sc-36024-SH and Nectin 3 shRNA (h) Lentiviral Particles: sc-36024-V.

Molecular Weight of Nectin 3: 83 kDa.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.