# AACT (71A1): sc-69751



The Power to Question

## **BACKGROUND**

The serine proteinase inhibitors (serpins) are a superfamily of proteins with a diverse set of functions, including the control of blood coagulation, complement activation, programmed cell death and development. The most abundant serpins in human plasma are  $\alpha\text{-1-antitrypsin}$  (AAT) and  $\alpha\text{-1-antichymotrypsin}$  (AACT). AACT (also called A1AC and SerpinA3) is a plasma protease inhibitor synthesized in the liver as a single glycopeptide chain. In humans, the normal serum level of AACT is about one-tenth that of AAT, with which it shares nucleic acid and protein sequence homology. Both are major acute phase reactants; their concentrations in plasma increase in response to trauma, surgery and infection. Elevated levels of AACT are widely, but not universally, reported in the cerebrospinal fluid and plasma of AD patients. Prostate-specific antigen (PSA) and its SDS-stable complex with AACT are in widespread use as markers for the diagnosis of prostate cancer. AACT deficiency may also be a possible cause of chronic liver disease.

## **REFERENCES**

- 1. Miyake, H., et al. 2001. Value of prostate specific antigen  $\alpha$ 1-antichymotrypsin complex for the detection of prostate cancer in patients with a PSA level of 4.1-10.0 ng/mL: comparison with PSA-related parameters. Int. J. Urol. 8: 589-593.
- 2. Kalsheker, N., et al. 2002. Gene regulation of the serine proteinase inhibitors  $\alpha$ 1-antitrypsin and  $\alpha$ 1-antichymotrypsin. Biochem. Soc. Trans. 30: 93-98.
- 3. Yoon, D., et al. 2002. Role of  $\alpha$ 1-antichymotrypsin deficiency in promoting cirrhosis in two siblings with heterozygous  $\alpha$ 1-antitrypsin deficiency phenotype SZ. Gut 50: 730-732.
- 4. Wang, X., et al. 2002. Distribution of plasma  $\alpha$ 1-antichymotrypsin levels in Alzheimer disease patients and controls and their genetic controls. Neurobiol. Aging 23: 377-382.
- 5. Hsieh, M.C., et al. 2002. Inhibition of prostate-specific antigen (PSA) by  $\alpha$ 1-antichymotrypsin: salt-dependent activation mediated by a conformational change. Biochem. 41: 2990-2997.
- 6. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 107280. World Wide Web URL: http://www.ncbi.nlm.nih.gov/omim/

## **CHROMOSOMAL LOCATION**

Genetic locus: SERPINA3 (human) mapping to 14g32.13.

## **SOURCE**

AACT (71A1) is a mouse monoclonal antibody raised against purified AACT of human origin.

## **PRODUCT**

Each vial contains 100  $\mu g \; lg G_1$  in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

#### **APPLICATIONS**

AACT (71A1) is recommended for detection of AACT of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)].

Suitable for use as control antibody for AACT siRNA (h): sc-40944, AACT shRNA Plasmid (h): sc-40944-SH and AACT shRNA (h) Lentiviral Particles: sc-40944-V.

Molecular Weight (predicted) of AACT: 48 kDa.


Molecular Weight (observed) of AACT: 65-76 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, T98G cell lysate: sc-2294 or SK-N-SH cell lysate: sc-2410.

#### **RECOMMENDED SECONDARY REAGENTS**

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-mouse IgG-HRP: sc-2005 (dilution range: 1:2000-1:32,000) or Cruz Marker™ compatible goat anti-mouse IgG-HRP: sc-2031 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-mouse IgG-FITC: sc-2010 (dilution range: 1:100-1:400) or goat anti-mouse IgG-TR: sc-2781 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



AACT (71A1): sc-69751. Western blot analysis of AACT purified from human plasma (**A**) and in human plasma (**B**)

#### **STORAGE**

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

# **PROTOCOLS**

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3801 Fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com