



Salmonella core antigen (6381): sc-69947

BACKGROUND

Salmonella is a genus of rod-shaped Gram-negative enterobacteria that causes typhoid fever, paratyphoid fever and foodborne illness. It is motile in nature and produces hydrogen sulfide. While salmonellosis is often considered to affect primarily the gastrointestinal tract, infection at other sites may occur, such as the peripheral or visceral arteries, with the abdominal aorta being the most frequent site of vascular infection. After a *Salmonella* infection, reactive arthritis (or Reiter's Syndrome) may develop. The *Salmonella* core antigen resides on *Salmonella enteritidis* D, *paratyphi* A, *typhimurium* B, *cholerae*, newport, *anatum* and *Selandia*.

REFERENCES

1. Cohen, J.I., Bartlett, J.A. and Corey, G.R. 1987. Extra-intestinal manifestations of *Salmonella* infections. *Medicine* 66: 349-388.
2. Jin, U.H., Chung, T.W., Lee, Y.C., Ha, S.D. and Kim, C.H. 2001. Molecular cloning and functional expression of the *rfaE* gene required for lipopolysaccharide biosynthesis in *Salmonella typhimurium*. *Glycoconj. J.* 18: 779-787.
3. Crich, D. and Li, H. 2002. Synthesis of the *Salmonella* type E₁ core trisaccharide as a probe for the generality of 1-(benzenesulfonyl) piperidine/triflic anhydride combination for glycosidic bond formation from thioglycosides. *J. Org. Chem.* 67: 4640-4646.
4. Graninger, M., Kneidinger, B., Bruno, K., Scheberl, A. and Messner, P. 2002. Homologs of the Rml enzymes from *Salmonella enterica* are responsible for dTDP-β-L-rhamnose biosynthesis in the gram-positive thermophile *Aneurinibacillus thermoaerophilus* DSM 10155. *Appl. Environ. Microbiol.* 68: 3708-3715.
5. Allende, D. and McIntosh, T.J. 2003. Lipopolysaccharides in bacterial membranes a plasma membranes in providing protection against melittin-induced bilayer lysis. *Biochemistry* 42: 1101-1108.
6. Abeyrathne, P.D., Daniels, C., Poon, K.K., Matewish, M.J. and Lam, J.S. 2005. *Funcni polysaccharide* to the core of *Pseudomonas aeruginosa* lipopolysaccharide. *J. Bacteriol.* 187: 3002-3012.
7. Frirdich, E., Bouwman, C., Vinogradov, E. and Whitfield, C. 2005. The role of galacturonic acid in outer membrane stability in *Klebsiella pneumoniae*. *J. Biol. Chem.* 280: 27604-27612.
8. Meng, F.P., Ding, J., Yu, Z.C., Han, Q.L., Guo, C.C., Liu, N. and Fan, D.M. 2005. Oral attenuated *Salmonella typhimurium* vaccine against MG7-Ag mimotope of gastric cancer. *World J. Gastroenterol.* 11: 1833-1836.
9. Woebking, B., Reuter, G., Shilling, R.A., Velamakanni, S., Shahi, S., Venter, H., Balakrishnan, L. and van Veen, H.W. 2005. Drug-lipid A interactions on the *Escherichia coli* ABC transporter MsbA. *J. Bacteriol.* 187: 6363-6369.

SOURCE

Salmonella core antigen (6381) is a mouse monoclonal antibody raised against *Salmonella* core antigen.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

Each vial contains 100 µg IgG_{2b} in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

Salmonella core antigen (6381) is recommended for detection of *Salmonella* O-serogroups A, B, C1, C2, D, E1, E3, E4, F, G1, G2 of *Salmonella* origin by immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.