

A20 (4H16): sc-69980

BACKGROUND

A20 is a Cys2/Cys2 zinc finger protein that is induced by a variety of inflammatory stimuli and regulates gene expression. Specifically, A20 is induced by tumor necrosis factor (TNF) and interleukin 1 (IL-1), and acts as a negative regulator of nuclear factor κ B (NF κ B) gene expression. By inhibiting NF κ B activation, A20 plays a critical role in terminating NF κ B responses to various stimuli. Although the C-terminal region of A20 contains seven zinc finger domains, only four of these domains are required for *in vitro* inhibition of TNF-induced NF κ B activation. A20 also interacts with several other proteins, such as TRAF2, TRAF6 and I κ B kinase (IKK) γ protein, and can thereby inhibit cell death. TXBP151, a novel A20-binding protein, may mediate the anti-apoptotic activity of A20. Involved in the negative feedback regulation of signal transduction, A20 and A20-binding proteins may be useful as novel therapeutic tools in the treatment of a variety of diseases.

REFERENCES

1. De Valck, D., et al. 1999. The zinc finger protein A20 interacts with a novel anti-apoptotic protein which is cleaved by specific caspases. *Oncogene* 18: 4182-4190.
2. Beyaert, R., et al. 2000. A20 and A20-binding proteins as cellular inhibitors of nuclear factor- κ B-dependent gene expression and apoptosis. *Biochem. Pharmacol.* 60: 1143-1151.
3. Van Huffel, S., et al. 2001. Identification of a novel A20-binding inhibitor of nuclear factor- κ B activation termed ABIN-2. *J. Biol. Chem.* 276: 30216-30223.
4. Lademann, U., et al. 2001. A20 zinc finger protein inhibits TNF-induced apoptosis and stress response early in the signaling cascades and independently of binding to TRAF2 or 14-3-3 proteins. *Cell Death Differ.* 8: 265-272.
5. Klinkenberg, M., et al. 2001. Functional redundancy of the zinc fingers of A20 for inhibition of NF κ B activation and protein-protein interactions. *FEBS Lett.* 498: 93-97.

CHROMOSOMAL LOCATION

Genetic locus: TNFAIP3 (human) mapping to 6q23.3; Tnfaip3 (mouse) mapping to 10 A3.

SOURCE

A20 (4H16) is a mouse monoclonal antibody raised against full length fusion A20 of human origin.

PRODUCT

Each vial contains 100 μ g IgG₁ in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS


A20 (4H16) is recommended for detection of A20 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)].

Suitable for use as control antibody for A20 siRNA (h): sc-37655, A20 siRNA (m): sc-37656, A20 shRNA Plasmid (h): sc-37655-SH, A20 shRNA Plasmid (m): sc-37656-SH, A20 shRNA (h) Lentiviral Particles: sc-37655-V and A20 shRNA (m) Lentiviral Particles: sc-37656-V.

Molecular Weight of A20: 90 kDa.

Positive Controls: Daudi cell lysate: sc-2415, Jurkat whole cell lysate: sc-2204 or U-937 cell lysate: sc-2239.

DATA



A20 (4H16): sc-69980. Western blot analysis of A20 expression in Jurkat (A), U-937 (B), Daudi (C) and HT-1080 (D) whole cell lysates.

SELECT PRODUCT CITATIONS

1. Shao, L., et al. 2013. A20 restricts wnt signaling in intestinal epithelial cells and suppresses colon carcinogenesis. *PLoS ONE* 8: e62223.
2. Mubarak, R.A., et al. 2018. Comparison of pro- and anti-inflammatory responses in paired human primary airway epithelial cells and alveolar macrophages. *Respir. Res.* 19: 126.
3. Nakamura, B.N., et al. 2018. A20 regulates canonical wnt-signaling through an interaction with RIPK4. *PLoS ONE* 13: e0195893.
4. Arora, H., et al. 2019. The ATP-binding cassette gene ABCF1 functions as an E2 ubiquitin-conjugating enzyme controlling macrophage polarization to dampen lethal septic shock. *Immunity* 50: 418-431.e6.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CONJUGATES

See **A20 (A-12): sc-166692** for A20 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.