

Carbonyl reductase 1 (I-15): sc-70215

BACKGROUND

Carbonyl reductase 1 and Carbonyl reductase 3 belong to the family of short-chain dehydrogenase/reductase proteins that play a role in metabolism throughout the body. Both proteins are monomeric carbonyl reductases that function to catalyze the NADPH-dependent reduction of various carbonyls (generally products of lipid peroxidation) to their corresponding alcohols. Carbonyl reductase 1 and Carbonyl reductase 3 share high sequence similarity at the amino acid level and are responsible for the metabolism of not only endogenous compounds, but of various pharmacological products as well. Genetic polymorphisms in both proteins result in individual variability at the level of drug metabolism. Defects in the genes encoding carbonyl reductase proteins have implications in cancer, diabetes and errors in metabolism.

REFERENCES

1. Watanabe, K., et al. 1999. Mapping of a novel human carbonyl reductase, CBR3 and ribosomal pseudogenes to human chromosome 21q22.2. *Genomics* 52: 95-100.
2. Terada, T., et al. 2001. Cloning and bacterial expression of monomeric short-chain dehydrogenase/reductase (Carbonyl reductase) from CHO-K1 cells. *Eur. J. Biochem.* 267: 6849-6857.
3. Olson, L.E., et al. 2003. Protection from doxorubicin-induced cardiac toxicity in mice with a null allele of Carbonyl reductase 1. *Cancer Res.* 63: 6602-6606.
4. Lakhman, S.S., et al. 2005. Functional significance of a natural allelic variant of human Carbonyl reductase 3 (CBR3). *Drug Metab. Dispos.* 33: 254-257.

CHROMOSOMAL LOCATION

Genetic locus: CBR1 (human) mapping to 21q22.12; Cbr1 (mouse) mapping to 16 C4.

SOURCE

Carbonyl reductase 1 (I-15) is an affinity purified rabbit polyclonal antibody raised against a peptide mapping within an internal region of Carbonyl reductase 1 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-70215 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

Carbonyl reductase 1 (I-15) is recommended for detection of Carbonyl reductase 1 of mouse and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Carbonyl reductase 1 (I-15) is also recommended for detection of Carbonyl reductase 1 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for Carbonyl reductase 1 siRNA (h): sc-72791, Carbonyl reductase 1 siRNA (m): sc-72792, Carbonyl reductase 1 shRNA Plasmid (h): sc-72791-SH, Carbonyl reductase 1 shRNA Plasmid (m): sc-72792-SH, Carbonyl reductase 1 shRNA (h) Lentiviral Particles: sc-72791-V and Carbonyl reductase 1 shRNA (m) Lentiviral Particles: sc-72792-V.

Molecular Weight of Carbonyl reductase 1: 30 kDa.

Positive Controls: Carbonyl reductase 1 (m): 293T Lysate: sc-118997, HeLa whole cell lysate: sc-2200 or NIH/3T3 whole cell lysate: sc-2210.

DATA



Carbonyl reductase 1 (I-15): sc-70215. Western blot analysis of Carbonyl reductase 1 expression in non-transfected: sc-117752 (A) and mouse Carbonyl reductase 1 transfected: sc-118997 (B) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **Carbonyl reductase 1 (B-11): sc-390554** or **Carbonyl reductase 1 (Z-8): sc-100518**, our highly recommended monoclonal alternatives to Carbonyl reductase 1 (I-15).