

B7RP-1 (3F206): sc-70398

BACKGROUND

T cell costimulatory molecule, inducible costimulator (ICOS)/B7-related protein-1 (B7RP-1, B7-H2, GL50, ICOS-L) is a ligand for the ICOS receptor that initiates T and B cell proliferation and cytokine secretion. B7RP-1 interactions play an essential role in T cell dependent B cell activation in peripheral lymphoid organs such as spleen and lymph nodes. B7RP-1 protein is present in myeloid leukocytes and by northern blot there are 2.4, 3.0 and 7.0 kb transcripts in brain, heart, kidney and liver, with lower expression in colon and thymus and a 1.1 kb transcript in leukocytes. Tumor necrosis factor α (TNF α), granulocyte-macrophage colony-stimulating factor (GM-CSF) and interleukin 4 (IL 4) enhance B7RP-1 expression. LPS induced up regulation of B7RP-1 is dependent on the MyD88 dependent signaling pathway.

REFERENCES

1. Yoshinaga, S.K., et al. 1999. T cell costimulation through B7RP-1 and ICOS. *Nature* 402: 827-832.
2. Yoshinaga, S.K., et al. 2000. Characterization of a new human B7 related protein: B7RP-1 is the ligand to the costimulatory protein ICOS. *Int. Immunol.* 12: 1439-1447.
3. Ling, V., et al. 2000. Cutting edge: identification of GL50, a novel B7 like protein that functionally binds to ICOS receptor. *J. Immunol.* 164: 1653-1657.
4. Richter, G., et al. 2001. Tumor receptor ligand on CD34+ progenitor cells during differentiation into antigen presenting cells. *J. Biol. Chem.* 276: 45686-45693.
5. Flesch, I.E. 2002. Inducible costimulator-ligand (ICOS-L). *J. Biol. Regul. Homeost. Agents* 16: 217-219.
6. Khayyamian, S., et al. 2002. ICOS-ligand, expressed on human endothelial cells, costimulates cytokine secretion by memory CD4+ T cells. *Proc. Natl. Acad. Sci. USA* 99: 6198-6203.
7. Iiyama, R., et al. 2003. The role of inducible costimulator (ICOS)/B7-related protein-1 (B7RP-1) interaction in the functional development of Peyer's patches. *Immunol. Lett.* 88: 63-70.
8. Wahl, P., et al. 2003. Interaction of B7RP-1 with ICOS negatively regulates antigen presentation by B cells. *Inflammation* 27: 191-200.
9. Gajewska, B.U., et al. 2005. B7RP-1 is not required for the generation of Th2 responses in a model of allergic airway inflammation but is essential for the induction of inhalation tolerance. *J. Immunol.* 174: 3000-3005.
10. Zhou, Z., et al. 2005. Antagonism between MyD88 and TRIF dependent signals in B7RP-1 up regulation. *Eur. J. Immunol.* 35: 1918-1927.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: ICOSLG (human) mapping to 21q22.3.

SOURCE

B7RP-1 (3F206) is a mouse monoclonal antibody raised against B7RP-1 of human origin.

PRODUCT

Each vial contains 100 μ g IgG_{2b} in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

B7RP-1 (3F206) is recommended for detection of B7RP-1 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)] and flow cytometry (1 μ g per 1 x 10⁶ cells).


Suitable for use as control antibody for B7RP-1 siRNA (h): sc-42768.

Molecular Weight of B7RP-1: 36 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-mouse IgG-HRP: sc-2005 (dilution range: 1:2000-1:32,000) or Cruz Marker™ compatible goat anti-mouse IgG-HRP: sc-2031 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

DATA



B7RP-1 (3F206): sc-70398. Western blot analysis of B7RP-1 expression in non-transfected: sc-117752 (A) and human B7RP-1 transfected: sc-116982 (B) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.