

# PKR (K-17): sc-707



The Power to Question

## BACKGROUND

Interferon-inducible RNA-dependent protein serine/threonine kinase, PKR, is variously designated in earlier literature as DA1, dsJ, PI kinase, p65, p67 or TIK for the mouse kinase; and p68, eIF-2 $\alpha$  protein kinase or p69 for the human kinase. The PKR kinase substrate is the  $\alpha$  subunit of protein synthesis initiation factor eIF-2. Phosphorylation of eIF-2 $\alpha$  on serine-51 results in inhibition of translation. Molecular cDNA clones have been isolated from both human and mouse cells. The serine/threonine kinase catalytic domains map to the carboxy-terminal half of the protein while the RNA-binding domains are located in the amino terminal region. Three kinds of regulation of PKR enzymatic activity have been described. These include transcriptional regulation in response to interferon, an autoregulatory mechanism controlling PKR expression at the level of translation and post-translational regulation by RNA-mediated autophosphorylation.

## CHROMOSOMAL LOCATION

Genetic locus: EIF2AK2 (human) mapping to 2p22.2.

## SOURCE

PKR (K-17) is an affinity purified rabbit polyclonal antibody raised against a peptide mapping at the C-terminus of PKR of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-707 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

PKR (K-17) is recommended for detection of PKR of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for PKR siRNA (h): sc-36263, PKR shRNA Plasmid (h): sc-36263-SH and PKR shRNA (h) Lentiviral Particles: sc-36263-V.

Molecular Weight of PKR: 68 kDa.

Positive Controls: A-431 whole cell lysate: sc-2201, HeLa whole cell lysate: sc-2200 or Daudi + IFN- $\alpha$  cell lysate: sc-2266.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.


## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.


## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## DATA



PKR (K-17): sc-707. Western blot analysis of PKR expression in IFN- $\alpha$ -induced Daudi whole cell lysate.



PKR (K-17): sc-707. Immunofluorescence staining of methanol-fixed Hep G2 cells showing cytoplasmic localization (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human urinary bladder tissue showing cytoplasmic staining of urothelial cells (B).

## SELECT PRODUCT CITATIONS

- Chou, J., et al. 1995. Association of a M(r) 90,000 phosphoprotein with protein kinase PKR in cells exhibiting enhanced phosphorylation of translation initiation factor eIF-2  $\alpha$  and premature shutoff of protein synthesis after infection with  $\gamma$  134.5-mutants of herpes simplex virus 1. Proc. Natl. Acad. Sci. USA 92: 10516-10520.
- Kang, J.I., et al. 2009. Cell culture-adaptive mutations in the NS5B gene of hepatitis C virus with delayed replication and reduced cytotoxicity. Virus Res. 144: 107-116.
- McAllister, C.S., et al. 2010. Mechanisms of protein kinase PKR-mediated amplification of  $\beta$  interferon induction by C protein-deficient measles virus. J. Virol. 84: 380-386.
- Muto, V., et al. 2011. Human papillomavirus type 16 E5 protein induces expression of beta interferon through interferon regulatory factor 1 in human keratinocytes. J. Virol. 85: 5070-5080.
- El Bougrini, J., et al. 2011. PML positively regulates interferon  $\gamma$  signaling. Biochimie 93: 389-398.
- Kang, J.I. and Ahn, B.Y. 2011. HCV-induced PKR activation is stimulated by the mitogen- and stress-activated protein kinase MSK2. Biochem. Biophys. Res. Commun. 407: 248-253.
- Blalock, W.L., et al. 2011. Multiple forms of PKR present in the nuclei of acute leukemia cells represent an active kinase that is responsive to stress. Leukemia 25: 236-245.
- Sanghvi, V.R., et al. 2011. Expression of interfering RNAs from an HIV-1 Tat-inducible chimeric promoter. Virus Res. 155: 106-111.



Try **PKR (B-10): sc-6282** or **PKR (H-12): sc-514626**, our highly recommended monoclonal alternatives to PKR (K-17). Also, for AC, HRP, FITC, PE, Alexa Fluor® 488 and Alexa Fluor® 647 conjugates, see **PKR (B-10): sc-6282**.