



CMV gB (5E258): sc-70943

BACKGROUND

Cytomegalovirus (CMV) is a member of the herpes virus group which includes herpes simplex virus types 1 and 2; Varicella Zoster Virus, which causes chicken pox; and Epstein Barr virus, which causes infectious mononucleosis. These viruses remain dormant within the body over a long period. In humans, CMV is known as HCMV or human herpesvirus 5 (HHV-5). HHV-5 causes only a brief mononeucleosis-like malaise in immunocompetent adults, but may cause severe illness or death in immunosuppressed individuals. CMV gB (CMV glycoprotein B) is an abundant virion envelope protein that is essential for the infectivity of CMV. CMV gB is also one of the most immunogenic virus-encoded proteins, and a significant fraction of virus neutralizing antibodies are directed at CMV gB.

REFERENCES

1. Alford, C.A., Stagno, S., Pass, R.F. and Britt, W.J. 1990. Congenital and perinatal Cytomegalovirus infections. *Rev. Infect. Dis.* 12: S745-753.
2. Rubin, R.H. 1990. Impact of Cytomegalovirus infection on organ transplant recipients. *Rev. Infect. Dis.* 12: S754-S766.
3. Toome, B.K., Bowers, K.E. and Scott, G.A. 1991. Diagnosis of cutaneous Cytomegalovirus infection: a review and report of a case. *J. Am. Acad. Dermatol.* 24: 860-867.
4. Kanj, S.S., Sharara, A.I., Clavien, P.A. and Hamilton, J.D. 1996. Cytomegalovirus infection following liver transplantation: review of the literature. *Clin. Infect. Dis.* 22: 537-549.
5. Boeckh, M. and Boivin, G. 1998. Quantitation of Cytomegalovirus: methodologic aspects and clinical applications. *Clin. Microbiol. Rev.* 11: 533-554.
6. Borchers, A.T., Perez, R., Kaysen, G., Ansari, A.A. and Gershwin, M.E. 1999. Role of Cytomegalovirus infection in mechanisms. *Transpl. Immunol.* 7: 75-82.
7. Drago, F., Aragone, M.G., Lugani, C. and Rebora, A. 2000. Cytomegalovirus infection in normal and immunocompromised humans. A review. *Dermatology* 200: 189-195.
8. Gaytant, M.A., Steegers, E.A., Semmekrot, B.A., Merkus, H.M. and Galama, J.M. 2002. Congenital Cytomegalovirus infection: review of the epidemiology and outcome. *Obstet. Gynecol. Surv.* 57: 245-256.
9. Fletcher, J.M., Vukmanovic-Stejic, M., Dunne, P.J., Birch, K.E., Cook, J.E., Jackson, S.E., Salmon, M., Rustin, M.H. and Akbar, A.N. 2005. Cytomegalovirus-specific CD4⁺ T cells in healthy carriers are continuously driven to replicative exhaustion. *J. Immunol.* 175: 8218-8225.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

SOURCE

CMV gB (5E258) is a mouse monoclonal antibody raised against CMV.

PRODUCT

Each vial contains 200 µg IgG₁ in 1.0 ml PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

CMV gB (5E258) is recommended for detection of CMV gB origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

Molecular Weight of CMV gB full length protein: 160 kDa.

Molecular Weight of CMV gB cleavage products: 55/110 kDa.

RESEARCH USE

For research use only, not for use in diagnostic procedures.