



CMV (1.B.214): sc-70944

BACKGROUND

Cytomegalovirus (CMV) is a member of the herpes virus group which includes herpes simplex virus types 1 and 2; Varicella Zoster Virus, which causes chicken pox; and Epstein Barr virus, which causes infectious mononucleosis. These viruses remain dormant within the body over a long period. In humans, CMV is known as HCMV or human herpesvirus 5 (HHV-5). It resides in body fluids, including urine, saliva, breast milk, blood, tears, semen and vaginal fluids. CMV especially targets salivary glands and may also be devastating or even fatal to fetuses. CMV infection can also be life threatening for patients who are immunocompromised, such as individuals with HIV or organ transplant recipients.

REFERENCES

1. Alford, C.A., Stagno, S., Pass, R.F. and Britt, W.J. 1990. Congenital and perinatal Cytomegalovirus infections. *Rev. Infect. Dis.* 12: S745-S753.
2. Rubin, R.H. 1990. Impact of Cytomegalovirus infection on organ transplant recipients. *Rev. Infect. Dis.* 12: S754-S766.
3. Toome, B.K., Bowers, K.E. and Scott, G.A. 1991. Diagnosis of cutaneous Cytomegalovirus infection: a review and report of a case. *J. Am. Acad. Dermatol.* 24: 860-867.
4. Kanj, S.S., Sharara, A.I., Clavien, P.A. and Hamilton, J.D. 1996. Cytomegalovirus infection following liver transplantation: review of the literature. *Clin. Infect. Dis.* 22: 537-549.
5. Boeckh, M. and Boivin, G. 1998. Quantitation of Cytomegalovirus: methodologic aspects and clinical applications. *Clin. Microbiol. Rev.* 11: 533-554.
6. Borchers, A.T., Perez, R., Kaysen, G., Ansari, A.A. and Gershwin, M.E. 1999. Role of Cytomegalovirus infection in mechanisms. *Transpl. Immunol.* 7: 75-82.
7. Drago, F., Aragone, M.G., Lugani, C. and Rebora, A. 2000. Cytomegalovirus infection in normal and immunocompromised humans. A review. *Dermatology* 200: 189-195.
8. Gaytant, M.A., Steegers, E.A., Semmekrot, B.A., Merkus, H.M. and Galama, J.M. 2002. Congenital Cytomegalovirus infection: review of the epidemiology and outcome. *Obstet. Gynecol. Surv.* 57: 245-256.
9. Fletcher, J.M., Vukmanovic-Stejic, M., Dunne, P.J., Birch, K.E., Cook, J.E., Jackson, S.E., Salmon, M., Rustin, M.H. and Akbar, A.N. 2005. Cytomegalovirus-specific CD4⁺ T cells in healthy carriers are continuously driven to replicative exhaustion. *J. Immunol.* 175: 8218-8225.

SOURCE

CMV (1.B.214) is a mouse monoclonal antibody raised against early proteins of CMV origin.

PRODUCT

Each vial contains 200 µg IgG_{2a} in 1.0 ml PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

CMV (1.B.214) is recommended for detection of a 65 kDa protein of CMV origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000).

Molecular Weight of CMV protein: 65 kDa.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

RESEARCH USE

For research use only, not for use in diagnostic procedures.