



EBV viral capsid antigen (1.B.265): sc-71029

BACKGROUND

Epstein-Barr virus (EBV), also designated human herpesvirus 4 (HHV-4), is a member of the herpesvirus family and is one of the most common human viruses, infecting about 90% of the population. EBV infects only B lymphocytes and, though often asymptomatic, it can cause infectious mononucleosis, a disease characterized by fatigue, fever, sore throat and muscle soreness. The linear genome of EBV circularizes once it enters the cell and exists there as an episome. EBV may play a role in the development of both Burkitt lymphoma, a disease in which a tumor can form on the mandible or maxilla, and nasopharyngeal carcinoma, a tumor found in the upper respiratory tract, most commonly in the nasopharynx. The viral capsid antigen (VCA) of EBV is used as a marker for screening for viral infection as well as nasopharyngeal carcinoma and many antigens from the viral capsid are used in diagnostic tests.

REFERENCES

- Luka, J., Miller, G., Jörnvall, H. and Pearson, G.R. 1986. Characterization of the restricted component of Epstein-Barr virus early antigens as a cytoplasmic filamentous protein. *J. Virol.* 58: 748-756.
- Goldschmidts, W.L., Ginsburg, M. and Pearson, G.R. 1989. Neutralization of Epstein-Barr virus-induced ribonucleotide reductase with antibody to the major restricted early antigen polypeptide. *Virology* 170: 330-333.
- Fan, J.A. 1989. Expression of the Epstein-Barr virus P150 viral capsid antigen in *Escherichia coli* for the use as antigen in diagnostic tests. *Zhongguo Yi Xue Ke Xue Yuan Xue Bao* 11: 381-387.
- Gorgievski-Hrisoho, M., Hinderer, W., Nebel-Schickel, H., Horn, J., Vornhagen, R., Sonneborn, H.H., Wolf, H. and Siegl, G. 1990. Serodiagnosis of infectious mononucleosis by using recombinant Epstein-Barr virus antigens and enzyme-linked immunosorbent assay technology. *J. Clin. Microbiol.* 28: 2305-2311.
- Tranchand-Bunel, D., Auriault, C., Diesis, E. and Gras-Masse, H. 1998. Detection of human antibodies using "convergent" combinatorial peptide libraries or "mixotopes" designed from a nonvariable antigen: application to the EBV viral capsid antigen p18. *J. Pept. Res.* 52: 495-508.
- Ruf, I.K., Rhyne, P.W., Yang, H., Borza, C.M., Hutt-Fletcher, L.M., Cleveland, J.L. and Sample, J.T. 1999. Epstein-Barr virus regulates c-Myc, apoptosis, and tumorigenicity in Burkitt lymphoma. *Mol. Cell. Biol.* 19: 1651-1660.
- Kantakamalakul, W., Chongkolwatana, C., Naksawat, P., Muangsomboon, S., Sukpanichnant, S., Chongvisal, S., Metheetrairat, C., Kositanont, U., and Puthavathana, P. 2000. Specific IgA antibody to Epstein-Barr viral capsid antigen: a better marker for screening nasopharyngeal carcinoma than EBV-DNA detection by polymerase chain reaction. *Asian Pac. J. Allergy Immunol.* 18: 221-226.
- Gan, Y.Y., Fones-Tan, A., Chan, S.H. and Gan, L.H. 2001. Epstein-Barr viral antigens used in the diagnosis of nasopharyngeal carcinoma. *J. Biomed. Sci.* 3: 159-169.
- Spender, L.C., Lucchesi, W., Bodelon, G., Bilancio, A., Karstegl, C.E., Asano, T., Dittrich-Breiholz, O., Kracht, M., Vanhaesebroeck, B. and Farrell, P.J. 2006. Cell target genes of Epstein-Barr virus transcription factor EBNA-2: induction of the p53 α regulatory subunit of PI3-kinase and its role in survival of EREB2.5 cells. *J. Gen. Virol.* 87: 2859-2867.

SOURCE

EBV viral capsid antigen (1.B.265) is a mouse monoclonal antibody raised against Epstein-Barr Virus.

PRODUCT

Each vial contains 100 μ g IgG₁ in 1.0 ml PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

EBV viral capsid antigen (1.B.265) is recommended for detection of EBV viral capsid antigen by immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Immunofluorescence: use goat anti-mouse IgG-FITC: sc-2010 (dilution range: 1:100-1:400) or goat anti-mouse IgG-TR: sc-2781 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.