

# FIV p24 (3H2128): sc-71120

## BACKGROUND

Feline immunodeficiency virus (FIV) is a lentivirus that affects domestic cats throughout the world. Five subtypes of FIV exist based on amino acid sequence differences in the viral envelope. The main mode of FIV transmission is through deep bite wounds, though it can also be spread via mucosal surfaces such as those in the mouth, rectum, and vagina. FIV is similar to Human Immunodeficiency Virus (HIV) in that it infects many cell types in its host, including CD4<sup>+</sup> and CD8<sup>+</sup> T lymphocytes, B lymphocytes, and macrophages. FIV eventually leads to collapse of the immune system because of the infection and exhaustion of the CD4<sup>+</sup> cells. FIV p120 and p24 are glycoproteins that localize to the envelope of the virus and are useful in diagnosis of the disease.

## REFERENCES

1. Pancino, G., Castelot, S. and Sonigo, P. 1995. Differences in feline immunodeficiency virus host cell range correlate with envelope fusogenic properties. *Virology* 206: 796-806.
2. Giannecchini, S., Di Fenza, A., D'Ursi, A.M., Matteucci, D., Rovero, P. and Bendinelli, M. 2003. Antiviral activity and conformational features of an octapeptide derived from the membrane-proximal ectodomain of the feline immunodeficiency virus transmembrane glycoprotein. *J. Virol.* 77: 3724-3733.
3. Lin, Y.C., Beck, Z., Morris, G.M., Olson, A.J. and Elder, J.H. 2003. Structural basis for distinctions between substrate and inhibitor specificities for feline immunodeficiency virus and human immunodeficiency virus proteases. *J. Virol.* 77: 6589-6600.
4. Shimojima, M., Nishimura, Y., Miyazawa, T., Tohya, Y. and Akashi, H. 2004. T cell subpopulations mediating inhibition of feline immunodeficiency virus replication in mucosally infected cats. *Microbes Infect.* 6: 265-271.
5. Reggeti, F. and Bienzle, D. 2004. Feline immunodeficiency virus subtypes A, B and C and intersubtype recombinants in Ontario, Canada. *J. Gen. Virol.* 85: 1843-1852.
6. Sinn, P.L., Burnight, E.R., Hickey, M.A., Blissard, G.W. and McCray, P.B. 2005. Persistent gene expression in mouse nasal epithelia following feline immunodeficiency virus-based vector gene transfer. *J. Virol.* 79: 12818-12827.
7. Dean, G.A., LaVoy, A., Yearley, J. and Stanton, C. 2006. Cytokine modulation of the innate immune response in feline immunodeficiency virus-infected cats. *J. Infect. Dis.* 193: 1520-1527.
8. Dunham, S.P., Bruce, J., MacKay, S., Golder, M., Jarrett, O. and Neil, J.C. 2006. Limited efficacy of an inactivated feline immunodeficiency virus vaccine. *Vet. Rec.* 158: 561-562.
9. Ryan, G., O'Flatharta, C., Callanan, J. and Mabruk, M.J. 2006. RNA *in situ* hybridization for the detection of feline immunodeficiency virus in infected cells. *Southeast Asian J. Trop. Med. Public Health* 37: 106-112.

## SOURCE

FIV p24 (3H2128) is a mouse monoclonal antibody raised against recombinant FIV p24.

## PRODUCT

Each vial contains 1.0 ml culture supernatant containing IgG<sub>1</sub> with < 0.1% sodium azide.

## APPLICATIONS

FIV p24 (3H2128) is recommended for detection of Gag p24 of FIV origin by Western Blotting (starting dilution to be determined by researcher, dilution range 1:10-1:200) and immunoprecipitation [10-20 µl per 100-500 µg of total protein (1 ml of cell lysate)].

Molecular Weight of FIV p24: 24 kDa.

## STORAGE

For immediate and continuous use, store at 4° C for up to one month. For sporadic use, freeze in working aliquots in order to avoid repeated freeze/thaw cycles. If turbidity is evident upon prolonged storage, clarify solution by centrifugation.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.