

Fyn (6A406): sc-71134

BACKGROUND

Src is the human homolog of the v-Src gene of the Rous sarcoma virus, also called avian sarcoma virus or ASV. Src was the first proto-oncogenic non-receptor tyrosine kinase characterized in human. By virtue of common structural motifs, the Src family is composed of nine members in vertebrates including Src, Yes, Fgr, Frk, Fyn, Lyn, Hck, Lck and Blk. Src-family kinases transduce signals that are involved in the control of a variety of cellular processes including proliferation, differentiation, motility and adhesion. Src-family kinases contain an amino-terminal cell membrane anchor followed by an SH3 domain and an SH2 domain involved in modular association and activation, respectively. Src-family kinases are normally maintained in an inactive state and can be activated transiently during cellular events such as mitosis. Different subcellular localizations of Src family kinases may be important for the regulation of specific cellular processes such as mitogenesis, cytoskeletal organization and membrane trafficking. Fyn and Lck kinases play a key role in T cell antigen receptor (TCR) signaling. The human Fyn gene maps to chromosome 6q21 and encodes a 537 amino acid protein.

REFERENCES

1. Sakaguchi, A.Y., et al. 1982. Organization of human proto-oncogenes. *Am. J. Hum. Genet.* 34: 175.
2. Hibbs, M.L., et al. 1997. Lyn, a Src-like tyrosine kinase. *Int. J. Biochem. Cell Biol.* 29: 397-400.
3. Williams, J.C., et al. 1998. Insights into Src kinase functions: structural comparisons. *Trends Biochem. Sci.* 23: 179-184.
4. Tatosyan, A.G., et al. 2000. Kinases of the Src family: structure and functions. *Biochemistry* 65: 49-58.
5. Bjorge, J.D., et al. 2000. Selected glimpses into the activation and function of Src kinase. *Oncogene* 19: 5620-5635.
6. Korade-Mirnic, Z., et al. 2000. Src kinase-mediated signaling in leukocytes. *J. Leukoc. Biol.* 68: 603-613.
7. Gilmore, E.S., et al. 2001. Src family kinases mediate epithelial Na⁺ channel inhibition by endothelin. *J. Biol. Chem.* 276: 42610-42617.

CHROMOSOMAL LOCATION

Genetic locus: FYN (human) mapping to 6q21; Fyn (mouse) mapping to 10 B1.

SOURCE

Fyn (6A406) is a mouse monoclonal antibody raised against amino acids 7-176 of Fyn of human origin.

PRODUCT

Each vial contains 100 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

Fyn (6A406) is recommended for detection of Fyn of mouse and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for Fyn siRNA (h): sc-29321, Fyn siRNA (m): sc-35425, Fyn shRNA Plasmid (h): sc-29321-SH, Fyn shRNA Plasmid (m): sc-35425-SH, Fyn shRNA (h) Lentiviral Particles: sc-29321-V and Fyn shRNA (m) Lentiviral Particles: sc-35425-V.


Molecular Weight of Fyn: 59 kDa.

Positive Controls: Fyn (h): 293T Lysate: sc-115186, HuT 78 whole cell lysate: sc-2208 or HeLa whole cell lysate: sc-2200.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-mouse IgG-HRP: sc-2005 (dilution range: 1:2000-1:32,000) or Cruz Marker™ compatible goat anti-mouse IgG-HRP: sc-2031 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-mouse IgG-FITC: sc-2010 (dilution range: 1:100-1:400) or goat anti-mouse IgG-TR: sc-2781 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



Fyn (6A406): sc-71134. Western blot analysis of Fyn expression in non-transfected 293T: sc-117752 (A), human Fyn transfected 293T: sc-115186 (B) and HeLa (C) whole cell lysates.



Fyn (6A406): sc-71134. Western blot analysis of Fyn expression in HuT 78 (A), Jurkat (B), HeLa (C) and M1 (D) whole cell lysates.

SELECT PRODUCT CITATIONS

1. Pullen, N.A., et al. 2012. Novel mechanism for FcεRI-mediated signal transducer and activator of transcription 5 (STAT5) tyrosine phosphorylation and the selective influence of STAT5B over mast cell cytokine production. *J. Biol. Chem.* 287: 2045-2054.
2. de la Puerta, M.L., et al. 2013. The autoimmunity risk variant LYP-W620 cooperates with CSK in the regulation of TCR signaling. *PLoS ONE* 8: e54569.

RESEARCH USE

For research use only, not for use in diagnostic procedures.