

CMV gB (1.B.223): sc-71228

BACKGROUND

Cytomegalovirus (CMV) is a member of the herpes virus group which includes herpes simplex virus types 1 and 2; Varicella Zoster Virus, which causes chicken pox; and Epstein Barr virus, which causes infectious mononucleosis. These viruses remain dormant within the body over a long period. In humans, CMV is known as HCMV or human herpesvirus 5 (HHV-5). HHV-5 causes only a brief mononeucleosis-like malaise in immunocompetent adults, but may cause severe illness or death in immunosuppressed individuals. CMV gB (CMV glycoprotein B) is an abundant virion envelope protein that is essential for the infectivity of CMV. CMV gB is also one of the most immunogenic virus-encoded proteins, and a significant fraction of virus neutralizing antibodies are directed at CMV gB.

REFERENCES

- Gönczöl, E., deTaisne, C., Hirka, G., Berencsi, K., Lin, W.C., Paoletti, E. and Plotkin, S. 1991. High expression of human cytomegalovirus (HCMV)-gB protein in cells infected with a vaccinia-gB recombinant: the importance of the gB protein in HCMV immunity. *Vaccine* 9: 631-637.
- Utz, U., Koenig, S., Coligan, J.E. and Biddison, W.E. 1992. Presentation of three different viral peptides, HTLV-1 Tax, HCMV gB, and influenza virus M1, is determined by common structural features of the HLA-A2.1 molecule. *J. Immunol.* 149: 214-221.
- Jarvis, M.A., Fish, K.N., Söderberg-Naucler, C., Streblov, D.N., Meyers, H.L., Thomas, G. and Nelson, J.A. 2002. Retrieval of human cytomegalovirus glycoprotein B from cell surface is not required for virus envelopment in astrocytoma cells. *J. Virol.* 76: 5147-5155.
- Carraro, E. and Granato, C.F. 2003. Single human cytomegalovirus gB genotype shed in multiple sites at the time of diagnosis in renal transplant recipients. *J. Med. Virol.* 70: 240-243.
- Crump, C.M., Hung, C.H., Thomas, L., Wan, L. and Thomas, G. 2003. Role of PACS-1 in trafficking of human cytomegalovirus glycoprotein B and virus production. *J. Virol.* 77: 11105-11113.
- Homman-Loudiyi, M., Hultenby, K., Britt, W. and Söderberg-Naucler, C. 2003. Envelopment of human cytomegalovirus occurs by budding into Golgi-derived vacuole compartments positive for gB, Rab 3, *trans*-Golgi network 46, and mannosidase II. *J. Virol.* 77: 3191-3203.
- Jarvis, M.A., Jones, T.R., Drummond, D.D., Smith, P.P., Britt, W.J., Nelson, J.A. and Baldick, C.J. 2003. Phosphorylation of human cytomegalovirus glycoprotein B (gB) at the acidic cluster casein kinase 2 site (Ser900) is required for localization of gB to the *trans*-Golgi network and efficient virus replication. *J. Virol.* 78: 285-293.
- Yue, Y., Zhou, S.S. and Barry, P.A. 2003. Antibody responses to rhesus cytomegalovirus glycoprotein B in naturally infected rhesus macaques. *J. Gen. Virol.* 84: 3371-3379.
- Heineman, T.C., Connolly, P., Hall, S.L. and Assefa, D. 2004. Conserved cytoplasmic and HCMV gB. *Virology* 328: 131-141.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

SOURCE

CMV gB (1.B.223) is a mouse monoclonal antibody raised against cells infected with CMV.

PRODUCT

Each vial contains 100 µg IgG₁ in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

CMV gB (1.B.223) is recommended for detection of gB of CMV by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

Molecular Weight of CMV gB full length protein: 160 kDa.

Molecular Weight of CMV gB cleavage products: 55/110 kDa.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.