

# Huntingtin (3H2242): sc-71287

## BACKGROUND

Huntingtin is a protein that contains a polyglutamine region. When the number of glutamine repeats exceeds 35, the gene encodes a version of Huntingtin that leads to Huntington's disease (HD). When the polyglutamine stretch is mutated, Huntingtin acts within the nucleus to induce neurodegeneration by a cell-specific apoptotic mechanism. Loss of Huntingtin activity is unlikely to be the cause of HD and it has been proposed that the expanded glutamine repeat region may induce an abnormal interaction between the mutant protein and other cellular proteins. Huntingtin interacts with a variety of proteins including HAP1, glyceraldehyde phosphate dehydrogenase (GAPDH) and HIP1.

## REFERENCES

1. The Huntington's Disease Collaborative Research Group. 1993. A novel gene containing a trinucleotide repeat that is expanded and unstable on Huntington's disease chromosomes. *Cell* 72: 971-983.
2. Ambrose, C.M., et al. 1994. Structure and expression of the Huntington's disease gene: evidence against simple inactivation due to an expanded CAG repeat. *Somat. Cell Mol. Genet.* 20: 27-38.
3. Albin, R.L. and Tagle, D.A. 1995. Genetics and molecular biology of Huntington's disease. *Trends Neurosci.* 18: 11-14.
4. Li, X.J., et al. 1995. A Huntingtin-associated protein enriched in brain with implications for pathology. *Nature* 378: 398-402.
5. Gusella, J.F., et al. 1996. Huntington's disease. *Cold Spring Harb. Symp. Quant. Biol.* 61: 615-626.
6. Burke, J.R., et al. 1996. Huntingtin and DRPLA proteins selectively interact with the enzyme GAPDH. *Nat. Med.* 2: 347-350.
7. Kalchman, M.A., et al. 1997. HIP1, a human homologue of *S. cerevisiae* Sla2p, interacts with membrane-associated Huntingtin in the brain. *Nat. Genet.* 16: 44-53.
8. Saudou, F., et al. 1998. Huntingtin acts in the nucleus to induce apoptosis but death does not correlate with the formation of intranuclear inclusions. *Cell* 95: 55-65.

## CHROMOSOMAL LOCATION

Genetic locus: HTT (human) mapping to 4p16.3; Htt (mouse) mapping to 5 B2.

## SOURCE

Huntingtin (3H2242) is a mouse monoclonal antibody raised against amino acids 997-1276 of Huntingtin of human origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

Huntingtin (3H2242) is recommended for detection of an epitope corresponding to the HDA region (amino acids 1171-1177) of Huntingtin of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for Huntingtin siRNA (h): sc-35617, Huntingtin siRNA (m): sc-35618, Huntingtin siRNA (r): sc-270267, Huntingtin shRNA Plasmid (h): sc-35617-SH, Huntingtin shRNA Plasmid (m): sc-35618-SH, Huntingtin shRNA Plasmid (r): sc-270267-SH, Huntingtin shRNA (h) Lentiviral Particles: sc-35617-V, Huntingtin shRNA (m) Lentiviral Particles: sc-35618-V and Huntingtin shRNA (r) Lentiviral Particles: sc-270267-V.


Molecular Weight of Huntingtin: 350 kDa.

Positive Controls: K-562 whole cell lysate: sc-2203, HeLa whole cell lysate: sc-2200 or Jurkat whole cell lysate: sc-2204.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## DATA



Huntingtin (3H2242): sc-71287. Western blot analysis of Huntingtin expression in HeLa whole cell lysate.



Huntingtin (3H2242): sc-71287. Western blot analysis of Huntingtin expression in K-562 whole cell lysate.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.