

ICAM-1 (2Q710): sc-71292

BACKGROUND

Cell adhesion molecules (CAMs) are a family of closely related cell surface glycoproteins involved in cell-cell interactions during growth and are thought to play important, yet separate, roles in embryogenesis and development. The intracellular adhesion molecule-1 (ICAM-1), also referred to as CD54, is an integral membrane protein of the immunoglobulin superfamily and recognizes the $\beta 2\alpha 1$ and $\beta 2\alpha M$ Integrins. ICAM-2 functions as a ligand for lymphocyte function-associated antigen-1 (LFA-1) and is involved in leukocyte adhesion. ICAM-3 is highly expressed on the surface of human eosinophils and, when bound to ligand, may inhibit eosinophil inflammatory responses and survival. ICAM-4, also known as LW glycoprotein, interacts with Integrins $\alpha L\beta 2$, $\alpha M\beta 2$, $\alpha 4\beta 1$, the αV family and $\alpha IIb\beta 3$, and selective binding to different integrins may be relevant to the pathology in a number of red blood cell associated diseases. Lastly, ICAM-5, expressed on telencephalic neurons, binds CD11a/CD18 and thus may act as an adhesion molecule for leukocyte binding in the central nervous system.

CHROMOSOMAL LOCATION

Genetic locus: ICAM1 (human) mapping to 19p13.2; Icam1 (mouse) mapping to 9 A3.

SOURCE

ICAM-1 (2Q710) is a mouse monoclonal antibody raised against an ICAM1-positive cell preparation.

PRODUCT

Each vial contains 200 μ g IgG κ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

ICAM-1 (2Q710) is recommended for detection of ICAM-1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and flow cytometry (1 μ g per 1 x 10⁶ cells).

Suitable for use as control antibody for ICAM-1 siRNA (h): sc-29354, ICAM-1 siRNA (m): sc-29355, ICAM-1 shRNA Plasmid (h): sc-29354-SH, ICAM-1 shRNA Plasmid (m): sc-29355-SH, ICAM-1 shRNA (h) Lentiviral Particles: sc-29354-V and ICAM-1 shRNA (m) Lentiviral Particles: sc-29355-V.


Molecular Weight of ICAM-1: 85-110 kDa.

Positive Controls: ICAM-1 (h): 293T Lysate: sc-176625, Raji whole cell lysate: sc-364236 or U266 whole cell lysate: sc-364800.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgG κ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



ICAM-1 (2Q710): sc-71292. Western blot analysis of ICAM-1 expression in Raji (A) and U266 (B) whole cell lysates.



ICAM-1 (2Q710): sc-71292. Western blot analysis of ICAM-1 expression in non-transfected: sc-117752 (A) and human ICAM-1 transfected: sc-176625 (B) 293T whole cell lysates.

SELECT PRODUCT CITATIONS

1. Tsutsumi, T., et al. 2010. Involvement of adhesion molecule in *in vitro* plaque-like formation of macrophages stimulated with *Aggregatibacter actinomycetemcomitans* lipopolysaccharide. J. Periodontal Res. 45: 550-556.
2. Qi, J., et al. 2017. Felodipine inhibits ox-LDL-induced reactive oxygen species production and inflammation in human umbilical vein endothelial cells. Mol. Med. Rep. 16: 4871-4878.
3. Li, X., et al. 2019. Down-regulation of Cx43 expression on PIH-HUVEC cells attenuates monocyte-endothelial adhesion. Thromb. Res. 179: 104-113.
4. Saadane, A., et al. 2020. Successful induction of diabetes in mice demonstrates no gender difference in development of early diabetic retinopathy. PLoS ONE 15: e0238727.

RESEARCH USE

For research use only, not for use in diagnostic procedures.



See **ICAM-1 (G-5): sc-8439** for ICAM-1 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.