

Integrin α X (2B2.36): sc-71453

BACKGROUND

Integrin α X (CD11c, leukocyte surface antigen p150,95, CR4, Axb2) is a type I transmembrane protein that traditionally combines with β 2 chain to form a leukocyte-specific integrin known as inactivated-C3b (iC3b) receptor 4 (CR4). Integrin α X/ β 2 shares similar properties of the α M/ β 2 Integrin in mediating adherence of neutrophils and monocytes to stimulated endothelial cells, and in phagocytosis of complement coated particles. Abnormal expression of Integrin α X is characteristic of hairy cell leukemia (HCL) and is dependent upon activation of proto-oncogenes Ras and Jun D. Proteins and DNA elements that influence transcription of Integrin α X include Sp1 and Sp1-like factors, AP-1 family, C/EBP, Oct-2 and PU.1. Integrin α X is present on monocyte derivative dendritic cells (DCs), macrophages and NK cells. Upon activation, DCs present in skin (Langerhans cells), lining of nose, lung, stomach, intestine and blood can migrate to lymphoid tissues and interact with T and B cells to initiate and shape the immune response.

REFERENCES

1. Nham, S.U. 1999. Characteristics of Fibrinogen binding to the domain of CD11c, an α subunit of p150,95. *Biochem. Biophys. Res. Commun.* 264: 630-634.
2. Binder, R.J., et al. 2000. Cutting edge: heat shock protein γ p96 induces maturation and migration of CD11c⁺ cells *in vivo*. *J. Immunol.* 165: 6029-6035.
3. Langegegen, H., et al. 2002. Human umbilical vein endothelial cells express complement receptor 1 (CD35) and complement receptor 4 (CD11c/CD18) *in vitro*. *Inflammation* 26: 103-110.
4. Nicolaou, F., et al. 2003. CD11c gene expression in hairy cell leukemia is dependent upon activation of the proto-oncogenes Ras and Jun D. *Blood* 101: 4033-4041.
5. Edwards, A.D., et al. 2003. Relationships among murine CD11c (high) dendritic cell subsets as revealed by baseline gene expression patterns. *J. Immunol.* 171: 47-60.
6. Paharkova-Vatchkova, V., et al. 2004. Estrogen preferentially promotes the differentiation of CD11c⁺ CD11b (intermediate) dendritic cells from bone marrow precursors. *J. Immunol.* 172: 1426-1436.
7. Scumpia, P.O., et al. 2005. CD11c⁺ dendritic cells are required for survival in murine polymicrobial sepsis. *J. Immunol.* 175: 3282-3286.

CHROMOSOMAL LOCATION

Genetic locus: ITGAX (human) mapping to 16p11.2.

SOURCE

Integrin α X (2B2.36) is a mouse monoclonal antibody raised against Integrin α X of human origin.

PRODUCT

Each vial contains 100 μ g IgG_{2a} in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

Integrin α X (2B2.36) is recommended for detection of Integrin α X of human origin by immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and flow cytometry (1 μ g per 1 x 10⁶ cells).

Suitable for use as control antibody for Integrin α X siRNA (h): sc-35695, Integrin α X shRNA Plasmid (h): sc-35695-SH and Integrin α X shRNA (h) Lentiviral Particles: sc-35695-V.

Molecular Weight of Integrin α X: 145 kDa.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.



See **Integrin α X (B-6): sc-46676** for Integrin α X antibody conjugates, including AC, HRP, FITC, PE, Alexa Fluor® 488 and Alexa Fluor® 647.