Integrin αX (3H986): sc-71456



The Power to Question

BACKGROUND

Integrin αX (CD11C, leukocyte surface antigen p150,95, CR4, Axb2) is a type 1 transmembrane protein that traditionally combines with $\beta 2$ chain to form a leukocyte-specific integrin known as inactivated-C3b (iC3b) receptor 4 (CR4). Integrin $\alpha X/\beta 2$ shares similar properties of the $\alpha M/\beta 2$ Integrin in mediating adherence of neutrophils and monocytes to stimulated endothelial cells and in phagocytosis of complement coated particles. Abnormal expression of Integrin αX is characteristic of hairy cell leukemia (HCL) and is dependent upon activation of proto-oncogenes Ras and JunD. Proteins and DNA elements that influence transcription of Integrin αX include Sp1 and Sp1-like factors, AP-1 family, C/EBP, Oct-2 and PU.1. Integrin αX is present on monocyte derivative dendritic cells (DCs), macrophages and NK cells. Upon activation, DCs present in skin (Langerhans cells), lining of nose, lung, stomach, intestine and blood can migrate to lymphoid tissues and interact with T and B cells to initiate and shape the immune response.

REFERENCES

- 1. Nham, S.U. 1999. Characteristics of Fibrinogen binding to the domain of CD11c, an α subunit of p150,95. Biochem. Biophys. Res. Commun. 264: 630-634.
- Binder, R.J., et al. 2000. Cutting edge: heat shock protein γp96 induces maturation and migration of CD11c+ cells in vivo. J. Immunol. 165: 6029-6035.
- Langeggen, H., et al. 2002. Human umbilical vein endothelial cells express complement receptor 1 (CD35) and complement receptor 4 (CD11c/CD18) in vitro. Inflammation 26: 103-110.
- Nicolaou, F., et al. 2003. CD11c gene expression in hairy cell leukemia is dependent upon activation of the proto-oncogenes Ras and JunD. Blood 101: 4033-4041.
- Edwards, A.D., et al. 2003. Relationships among murine CD11c (high) dendritic cell subsets as revealed by baseline gene expression patterns. J. Immunol. 171: 47-60.
- Paharkova-Vatchkova, V., et al. 2004. Estrogen preferentially promotes the differentiation of CD11c+ CD11b (intermediate) dendritic cells from bone marrow precursors. J. Immunol. 172: 1426-1436.
- 7. Scumpia, P.O., et al. 2005. CD11c+ dendritic cells are required for survival in murine polymicrobial sepsis. J. Immunol. 175: 3282-3286.
- 8. Sundquist, M., et al. 2005. TNF α -dependent and -independent maturation of dendritic cells and recruited CD11c (int) CD11b⁺ cells during oral salmonella infection. J. Immunol. 175: 3287-3298.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: Itgax (mouse) mapping to 7 F3.

SOURCE

Integrin αX (3H986) is a Armenian hamster monoclonal antibody raised against mouse spleen dendritic cells.

PRODUCT

Each vial contains 200 μg lgG in 1.0 ml PBS with < 0.1% sodium azide and 0.1% gelatin.

Integrin α X (3H986) is available conjugated to either phycoerythrin (sc-71456 PE) or fluorescein (sc-71456 FITC), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM.

APPLICATIONS

Integrin αX (3H986) is recommended for detection of Integrin αX of mouse origin by immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and flow cytometry (1 μg per 1 x 10⁶ cells).

Suitable for use as control antibody for Integrin α X siRNA (m): sc-35696, Integrin α X shRNA Plasmid (m): sc-35696-SH and Integrin α X shRNA (m) Lentiviral Particles: sc-35696-V.

Molecular Weight of Integrin α X: 145 kDa.

RESEARCH USE

For research use only, not for use in diagnostic procedures.



See Integrin α X (D-8): sc-398708 for Integrin α X antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3801 Fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com