

Ku70 (3F246): sc-71469

BACKGROUND

The Ku protein is localized in the nucleus and is composed of subunits referred to as Ku70 (p70) and Ku86 (p86) which is also known by the synonym Ku80 or (p80). Ku was first described as an autoantigen to which antibodies were produced in a patient with scleroderma polymyositis overlap syndrome, and was later found in the sera of patients with other rheumatic diseases. Both subunits of the Ku protein have been cloned, and a number of functions have been proposed for Ku, including cell signaling, DNA replication and transcriptional activation. Ku is involved in Pol II-directed transcription by virtue of its DNA binding activity, serving as the regulatory component of the DNA-associated protein kinase that phosphorylates Pol II and transcription factor Sp. Ku proteins also activate transcription from the U1 small nuclear RNA and the human transferrin receptor gene promoters. A Ku-related protein designated the enhancer 1 binding factor (E1BF), composed of two subunits, has been identified as a positive regulator of RNA polymerase I transcription initiation.

REFERENCES

- Mimori, T., et al. 1981. Characterization of a high molecular weight acidic nuclear protein recognized by autoantibodies in sera from patients with polymyositis-scleroderma overlap. *J. Clin. Invest.* 68: 611-620.
- Mimori, T., et al. 1986. Characterization of the DNA-binding protein antigen Ku recognized by autoantibodies from patients with rheumatic disorders. *J. Biol. Chem.* 261: 2274-2278.

CHROMOSOMAL LOCATION

Genetic locus: XRCC6 (human) mapping to 22q13.2; Xrcc6 (mouse) mapping to 15 E1.

SOURCE

Ku70 (3F246) is a mouse monoclonal antibody raised against PSE1-PL placental cells of human origin.

PRODUCT

Each vial contains 200 µg IgG_{2b} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

Ku70 (3F246) is recommended for detection of Ku70 of mouse, rat, human and *Xenopus laevis* origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for Ku70 siRNA (h): sc-29383, Ku70 siRNA (m): sc-35764, Ku70 shRNA Plasmid (h): sc-29383-SH, Ku70 shRNA Plasmid (m): sc-35764-SH, Ku70 shRNA (h) Lentiviral Particles: sc-29383-V and Ku70 shRNA (m) Lentiviral Particles: sc-35764-V.


Molecular Weight of Ku70: 70 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, Jurkat whole cell lysate: sc-2204 or HEL 92.1.7 cell lysate: sc-2270.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



Ku70 (3F246): sc-71469. Western blot analysis of Ku70 expression in HeLa (A), Jurkat (B) and HEL 92.1.7 (C) whole cell lysates.



Ku70 (3F246): sc-71469. Immunoperoxidase staining of formalin fixed, paraffin-embedded human testis tissue showing nuclear staining of cells in seminiferous ducts and Leydig cells (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human skin tissue showing nuclear staining of keratinocytes, fibroblasts, Langerhans cells and melanocytes (B).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.



See **Ku70 (E-5): sc-17789** for Ku70 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.