

NFκB p65 (4H212): sc-71677

BACKGROUND

Proteins encoded by the v-Rel viral oncogene and its cellular homolog, c-Rel, are members of a family of transcription factors that include the two subunits of the transcription factor NFκB (p50 and p65) and the *Drosophila* maternal morphogen, dorsal. Both proteins specifically bind to DNA sequences that are the same or slight variations of the 10 bp κB sequence in the immunoglobulin κ light chain enhancer. This same sequence is also present in a number of other cellular and viral enhancers. The DNA binding activity of NFκB is activated and NFκB is subsequently transported from the cytoplasm to the nucleus in cells exposed to mitogens or growth factors. cDNAs encoding precursors for two distinct proteins of the same size have been described, designated p105 and p100. The p105 precursor contains p50 at its N-terminus and a C-terminal region that when expressed as a separate molecule, designated pdl, binds to p50 and regulates its activity.

CHROMOSOMAL LOCATION

Genetic locus: RELA (human) mapping to 11q13.1; Rela (mouse) mapping to 19 A.

SOURCE

NFκB p65 (4H212) is a mouse monoclonal antibody raised against amino acids 526-539 of NFκB p65 of human origin.

PRODUCT

Each vial contains 100 μg IgG₁ in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

NFκB p65 (4H211) is recommended for detection of NFκB p65 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and immunoprecipitation [1-2 μg per 100-500 μg of total protein (1 ml of cell lysate)].

Suitable for use as control antibody for NFκB p65 siRNA (h): sc-29410, NFκB p65 siRNA (m): sc-29411, NFκB p65 shRNA Plasmid (h): sc-29410-SH, NFκB p65 shRNA Plasmid (m): sc-29411-SH, NFκB p65 shRNA (h) Lentiviral Particles: sc-29410-V and NFκB p65 shRNA (m) Lentiviral Particles: sc-29411-V.

Molecular Weight of NFκB p65: 65 kDa.

Positive Controls: NFκB p65 (m): 293T Lysate: sc-122027, Jurkat whole cell lysate: sc-2204 or K-562 whole cell lysate: sc-2203.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.


PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.


RESEARCH USE

For research use only, not for use in diagnostic procedures.

DATA



NFκB p65 (4H212): sc-71677. Western blot analysis of NFκB p65 expression in K-562 (A), Jurkat (B), HUV-EC-C (C) and HeLa (D) whole cell lysates.



NFκB p65 (4H212): sc-71677. Western blot analysis of NFκB p65 expression in non-transfected: sc-117752 (A) and mouse NFκB p65 transfected: sc-122027 (B) 293T whole cell lysates.

SELECT PRODUCT CITATIONS

- Wu, C.M., et al. 2011. IGF-I enhances α5β1 integrin expression and cell motility in human chondrosarcoma cells. *J. Cell. Physiol.* 226: 3270-3277.
- Fu, X.D., et al. 2012. Dydrogesterone exerts endothelial anti-inflammatory actions decreasing expression of leukocyte adhesion molecules. *Mol. Hum. Reprod.* 18: 44-51.
- Jia, C.H., et al. 2013. IKK-β mediates hydrogen peroxide induced cell death through p85 S6K1. *Cell Death Differ.* 20: 248-258.
- Eisfeld, A.K., et al. 2014. Intronic miR-3151 within BAALC drives leukemogenesis by deregulating the TP53 pathway. *Sci. Signal.* 7: ra36.
- Pandit, H., et al. 2015. Manganese superoxide dismutase expression is negatively associated with microRNA-301a in human pancreatic ductal adenocarcinoma. *Cancer Gene Ther.* 22: 481-486.
- Gai, J.W., et al. 2018. Expression of CD74 in bladder cancer and its suppression in association with cancer proliferation, invasion and angiogenesis in HT-1376 cells. *Oncol. Lett.* 15: 7631-7638.
- Zhang, L., et al. 2019. Silence of lncRNA CHRFB protects H9c2 cells against lipopolysaccharide-induced injury via up-regulating microRNA-221. *Exp. Mol. Pathol.* 107: 43-50.
- Li, W., et al. 2019. MicroRNA-451 relieves inflammation in cerebral ischemia-reperfusion via the Toll-like receptor 4/MyD88/NFκB signaling pathway. *Mol. Med. Rep.* 20: 3043-3054.
- Wang, R., et al. 2019. Ginsenoside metabolite compound-K regulates macrophage function through inhibition of β-arrestin2. *Biomed. Pharmacother.* 115: 108909.

CONJUGATES

See **NFκB p65 (F-6): sc-8008** for NFκB p65 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor[®] 488, 546, 594, 647, 680 and 790.