

# NGFR p75 (0.N.469) sc-71695

## BACKGROUND

The Trk oncogene encodes a membrane-spanning protein tyrosine kinase, gp140Trk, whose expression is restricted *in vivo* to neurons of the sensory spinal and cranial ganglia of neural crest origin. Nerve growth factor (NGF) stimulates tyrosine phosphorylation of Trk A in neural cell lines and in embryonic dorsal root ganglia. Tyrosine phosphorylation of Trk by NGF is rapid, specific and occurs with picomolar quantities of factor, indicating that the response is mediated by physiological amounts of NGF, suggesting that Trk A participates in the primary signal transduction mechanism of NGF. An additional component of the Trk A receptor complex, NGFR p75, binds to the neurotrophic factors with low affinity but is required for efficient signaling. NGFR p75 accelerates Trk A activation and may recruit downstream effector molecules to the liganded complex.

## REFERENCES

1. Martin-Zanca, D., et al. 1986. A human oncogene formed by the fusion of truncated Tropomyosin and protein tyrosine kinase sequences. *Nature* 319: 743-748.
2. Reinach, F.C., et al. 1986. Tissue-specific expression of the human Tropomyosin gene involved in the generation of the Trk oncogene. *Nature* 322: 648-650.
3. Martin-Zanca, D., et al. 1989. Molecular and biochemical characterization of the human Trk proto-oncogene. *Mol. Cell. Biol.* 9: 24-33.

## CHROMOSOMAL LOCATION

Genetic locus: NGFR (human) mapping to 17q21.33.

## SOURCE

NGFR p75 (0.N.469) is a mouse monoclonal antibody raised against NGFR p75 from A875 melanoma cells of human origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## APPLICATIONS

NGFR p75 (0.N.469) is recommended for detection of NGFR p75 of human and feline origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and flow cytometry (1 µg per 1 x 10<sup>6</sup> cells); non cross-reactive with mouse or rat NGFR p75.

Suitable for use as control antibody for NGFR p75 siRNA (h): sc-36058, NGFR p75 shRNA Plasmid (h): sc-36058-SH and NGFR p75 shRNA (h) Lentiviral Particles: sc-36058-V.


Molecular Weight of NGFR p75: 75 kDa.

Positive Controls: SK-N-MC cell lysate: sc-2237.

## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## DATA



NGFR p75 (0.N.469): sc-71695. Western blot analysis of NGFR p75 expression in SK-N-MC whole cell lysate.

## SELECT PRODUCT CITATIONS

1. Dalley, A.J., et al. 2012. Organotypic culture of normal, dysplastic and squamous cell carcinoma-derived oral cell lines reveals loss of spatial regulation of CD44 and p75(NTR) in malignancy. *J. Oral Pathol. Med.* 42: 37-46.
2. Abdulmajeed, A.A., et al. 2013. Putative cancer stem cell marker expression in oral epithelial dysplasia and squamous cell carcinoma. *J. Oral. Pathol. Med.* 42: 755-760.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.



See **NGFR p75 (B-1): sc-271708** for NGFR p75 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.