Nop5p (4i247): sc-71717



The Power to Question

BACKGROUND

Nop1p (nucleolar protein 1) is a phylogenetically conserved protein essential for efficient processing of pre-rRNA through its association with a class of small nucleolar RNAs during ribosomal biogenesis. Small nucleolar RNAs (snoRNAs) are associated in ribonucleoprotein particles localized to the nucleolus (snoRNPs). Nop1p is structurally and functionally homologous to vertebrate Fibrillarin and is essential for viability. The *Saccharomyces cerevisiae* NOP1 gene encodes a protein resembling the dense fibrillar region of mammalian nucleoli. Nop5p functions with Nop1p in the execution of early pre-rRNA processing steps that lead to formation of 18 S rRNA. In *Archaea*, Fibrillarin and Nop5p comprise the core complex of box C/D snoRNAs, which are responsible for site-specific 2'-hydroxyl methylation of ribosomal and transfer RNAs.

REFERENCES

- Wu, P., et al. 1998. Nop5p is a small nucleolar ribonucleoprotein component required for pre-18 S rRNA processing in yeast. J. Biol. Chem. 273: 16453-16463.
- Lafontaine, D.L., et al. 2000. Synthesis and assembly of the box C/D small nucleolar RNPs. Mol. Cell. Biol. 20: 2650-2659.
- Nelson, S.A., et al. 2000. Multiple growth factor induction of a murine early response gene that complements a lethal defect in yeast ribosome biogenesis. J. Biol. Chem. 275: 13835-13841.
- Verheggen, C., et al. 2001. Box C/D small nucleolar RNA trafficking involves small nucleolar RNP proteins, nucleolar factors and a novel nuclear domain. EMBO J. 20: 5480-5490.
- Aittaleb, M., et al. 2003. Structure and function of archaeal box C/D sRNP core proteins. Nat. Struct. Biol. 10: 256-263.
- Rashid, R., et al. 2003. Functional requirement for symmetric assembly of archaeal box C/D small ribonucleoprotein particles. J. Mol. Biol. 333: 295-306.
- 7. Bortolin, M.L., et al. 2003. *In vitro* RNP assembly and methylation guide activity of an unusual box C/D RNA, *cis*-acting archaeal pre-tRNA (Trp). Nucleic Acids Res. 31: 6524-6535.
- Aittaleb, M., et al. 2004. Structural and thermodynamic evidence for a stabilizing role of Nop5p in S-adenosyl-L-methionine binding to Fibrillarin. J. Biol. Chem. 279: 41822-41829.
- Hardin, J.W., et al. 2006. The bipartite architecture of the sRNA in an archaeal box C/D complex is a primary determinant of specificity. Nucleic Acids Res. 34: 5039-5051.

SOURCE

Nop5p (4i247) is a mouse monoclonal antibody raised against a yeast nucleolar preparation.

PRODUCT

Each vial contains 500 μl culture supernatant containing IgG in PBS with < 0.1% sodium azide.

APPLICATIONS

Nop5p (4i247) is recommended for detection of Nop5p of *S. cerevisiae* origin by Western Blotting (starting dilution to be determined by researcher, dilution range 1:1000-1:5000), immunoprecipitation [2-4 μ l per 100-500 μ g of total protein (1 ml of cell lysate)] and immunofluorescence (starting dilution to be determined by researcher, dilution range 1:5000-1:25000).

Molecular Weight of Nop5p: 58 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-mouse IgG-HRP: sc-2005 (dilution range: 1:2000-1:32,000) or Cruz Marker™ compatible goat anti-mouse IgG-HRP: sc-2031 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-mouse IgG-FITC: sc-2010 (dilution range: 1:100-1:400) or goat anti-mouse IgG-TR: sc-2781 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

For immediate and continuous use, store at 4° C for up to one month. For sporadic use, freeze in working aliquots in order to avoid repeated freeze/thaw cycles. If turbidity is evident upon prolonged storage, clarify solution by centrifugation.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3801 fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com