

# ORC1 (3H2792): sc-71752

## BACKGROUND

The initiation of DNA replication is a multi-step process that depends on the formation of pre-replication complexes, which trigger initiation. Among the proteins required for establishing these complexes are the origin recognition complex (ORC) proteins. ORC proteins bind specifically to origins of replication where they serve as scaffold for the assembly of additional initiation factors. Human ORC subunits 1-6 are expressed in the nucleus of proliferating cells and tissues, such as the testis. ORC1 and ORC2 are both expressed at equivalent concentrations throughout the cell cycle; however, only ORC2 remains stably bound to chromatin. ORC4 and ORC6 are also expressed constantly throughout the cell cycle. ORC2, ORC3, ORC4 and ORC5 form a core complex upon which ORC6 and ORC1 assemble. The formation of this core complex suggests that ORC proteins play a crucial role in the G<sub>1</sub>-S transition in mammalian cells.

## REFERENCES

1. Quintana, D.G., Hou, Z., Thome, K.C., Hendricks, M., Saha, P. and Dutta, A. 1997. Identification of the HsORC4, a member of the human origin of replication recognition complex. *J. Biol. Chem.* 272: 28247-28251.
2. Mendez, J. and Stillman, B. 2000. Chromatin association of human origin recognition complex, Cdc6, and minichromosome maintenance proteins during the cell cycle: assembly of prereplication complexes in late mitosis. *Mol. Cell. Biol.* 20: 8602-8612.
3. Dhar, S.K. and Dutta, A. 2000. Identification and characterization of the human ORC6 homolog. *J. Biol. Chem.* 275: 34983-34988.
4. Thome, K.C., Dhar, S.K., Quintana, D.G., Delmolino, L., Shahsafaei, A. and Dutta, A. 2000. Subsets of human origin recognition complex (ORC) subunits are expressed in non-proliferating cells and associate with non-ORC proteins. *J. Biol. Chem.* 275: 35233-35241.
5. Kreitz, S., Ritzi, M., Baack, M. and Knippers, R. 2000. The human origin recognition complex protein 1 dissociates from chromatin during S phase in HeLa cells. *J. Biol. Chem.* 276: 6337-6342.
6. Natale, D.A., Li, C.J., Sun, W.H. and DePamphilis, M.L. 2000. Selective instability of ORC1 protein accounts for the absence of functional origin recognition complexes during the M-G<sub>1</sub> transition in mammals. *EMBO J.* 19: 2728-2738.
7. Vashee, S., Simancek, P., Challberg, M.D. and Kelly, T.J. 2001. Assembly of the human origin recognition complex. *J. Biol. Chem.* 276: 26666-26673.

## CHROMOSOMAL LOCATION

Genetic locus: ORC1L (human) mapping to 1p32.3.

## SOURCE

ORC1 (3H2792) is a mouse monoclonal antibody raised against recombinant ORC1 of human origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## APPLICATIONS

ORC1 (3H2792) is recommended for detection of ORC1 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)].

Suitable for use as control antibody for ORC1 siRNA (h): sc-38151, ORC1 shRNA Plasmid (h): sc-38151-SH and ORC1 shRNA (h) Lentiviral Particles: sc-38151-V.

Molecular Weight of ORC1: 120 kDa.

Positive Controls: Ramos nuclear extract: sc-2153.

## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml).

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.