

# p-EGFR (6D371): sc-71777

## BACKGROUND

Epidermal growth factor mediates its effects on cell growth through its interaction with a cell surface glycoprotein designated the EGF receptor. Binding of EGF or TGF $\alpha$  to the EGF receptor activates tyrosine-specific protein kinase activity intrinsic to the EGF receptor. The carboxy-terminal tyrosine residues on EGFR, Tyr 1068 and Tyr 1173 are the major sites of autophosphorylation, which occurs as a result of EGF binding. Once activated, EGFR mediates the binding of the phosphotyrosine binding (PTB) domain of GRB2 through direct interactions with Tyr 1068 and Tyr 1086 and through indirect interactions with Tyr 1173 in the Ras signaling pathway. Tyr 1173 of EGFR also functions as a kinase substrate. Phosphorylation of Tyr 992, Tyr 1068 and Tyr 1086 is required for conformational change in the C-terminal tail of the EGF receptor.

## REFERENCES

1. Reynolds, F.H., Jr., Todaro, G.J., Fryling, C. and Stephenson, J.R. 1981. Human transforming growth factors induces tyrosine phosphorylation of EGF receptors. *Nature* 292: 259-262.
2. Hunter, T. 1984. The epidermal growth factor receptor gene and its product. *Nature* 311: 414-416.
3. Batzer, A.G., Rotin, D., Urena, J.M., Skolnik, E.Y. and Schlessinger, J. 1994. Hierarchy of binding site for GRB2 and Shc on the epidermal growth factor receptor. *Mol. Cell. Biol.* 14: 5192-5201.
4. Ward, C.W., Gough, K.H., Rashke, M., Wan, S.S., Tribbick, G. and Wang, J.X. 1996. Systematic mapping of potential binding sites for Shc and GRB2 SH2 domains on Insulin receptor substrate-1 and the receptors for Insulin, epidermal growth factor, platelet-derived growth factor, and fibroblast growth factor. *J. Biol. Chem.* 271: 5603-5609.
5. Rojas, M., Yao, S. and Lin, Y. Z. 1996. Controlling epidermal growth factor (EGF)-stimulated Ras activation in intact cells by a cell-permeable peptide mimicking phosphorylated EGF receptor. *J. Biol. Chem.* 271: 27456-27461.
6. Wright, J.D., Reuter, C.W. and Weber, M.J. 1996. Identification of sites on epidermal growth factor receptors which are phosphorylated by pp60src *in vitro*. *Biochim. Biophys. Acta* 1312: 85-93.
7. Sakaguchi, K., Okabayashi, Y., Kido, Y., Kimura, S., Matsumura, Y., Inushima, K. and Kasuga, M. 1998. Shc phosphotyrosine-binding domain dominantly interacts with epidermal growth factor receptors and mediates Ras activation in intact cells. *Mol. Endocrinol.* 12: 536-543.
8. Bishayee, A., Beguinot, L. and Bishayee, S. 1999. Phosphorylation of Tyrosine 992, 1068 and 1086 is required for conformational change of the human epidermal growth factor receptor C-terminal tail. *Mol. Biol. Cell* 10: 525-536.

## CHROMOSOMAL LOCATION

Genetic locus: EGFR (human) mapping to 7p11.2; Egfr (mouse) mapping to 11 A2.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## SOURCE

p-EGFR (6D371) is a mouse monoclonal antibody raised against a short amino acid sequence containing phosphorylated raised against an EGFR phosphopeptide of human origin of EGFR of origin.

## PRODUCT

Each vial contains 50  $\mu$ g IgG<sub>1</sub> in 0.5 ml of PBS with < 0.1% sodium azide, 0.1% gelatin, PEG and sucrose.

## APPLICATIONS

p-EGFR (6D371) is recommended for detection of Tyr 845 phosphorylated EGFR of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000); non cross-reactive with the non-phosphorylated EGFR nor with unrelated Tyrosine-phosphorylated proteins.

Suitable for use as control antibody for EGFR siRNA (h): sc-29301, EGFR siRNA (m): sc-29302, EGFR siRNA (r): sc-108050, EGFR shRNA Plasmid (h): sc-29301-SH, EGFR shRNA Plasmid (m): sc-29302-SH, EGFR shRNA Plasmid (r): sc-108050-SH, EGFR shRNA (h) Lentiviral Particles: sc-29301-V, EGFR shRNA (m) Lentiviral Particles: sc-29302-V and EGFR shRNA (r) Lentiviral Particles: sc-108050-V.

Molecular Weight of p-EGFR: 170 kDa.

Positive Controls: A-431+EGF whole cell lysate: sc-2202, SK-OV-3 whole cell lysate: sc-364229 or SK-N-SH cell lysate: sc-2410.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.



See **p-EGFR (A-10): sc-373746** for p-EGFR antibody conjugates, including AC, HRP, FITC, PE, Alexa Fluor® 488 and Alexa Fluor® 647.