# SANTA CRUZ BIOTECHNOLOGY, INC.

# Lamin A/C (346): sc-7293



# BACKGROUND

A unique family of cysteine proteases has been described that differs in sequence, structure and substrate specificity from any previously described protease family. This family, termed CED-3/ICE, is comprised of ICE, CPP32, ICH-1/Nedd-2, Tx, Mch2, Mch3 (ICE-LAP3 or CMH-1), Mch4 and ICE-LAP6. CED-3/ICE family members function as key components of the apoptotic machinery and act to destroy specific target proteins which are critical to cellular longevity. Nuclear lamins are critical to maintaining the integrity of the nuclear envelope and cellular morphology. The nuclear Lamin A is cleaved by Mch2, but not CPP32. Nuclear Lamin B is fragmented as a consequence of apoptosis by an unidentified member of the ICE family. Lamin C is a splice variant of Lamin A, differing only at the carboxy-terminus. Lamins A and C are identical for the first 566 amino acids, with Lamin C differing only in six unique carboxy-terminal amino acids.

## **CHROMOSOMAL LOCATION**

Genetic locus: LMNA (human) mapping to 1q22; Lmna (mouse) mapping to 3 F1.

#### SOURCE

Lamin A/C (346) is a mouse monoclonal antibody raised against nuclear lamin fraction of rat liver origin.

# PRODUCT

Each vial contains 200  $\mu g$  lgM kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Lamin A/C (346) is available conjugated to agarose (sc-7293 AC), 500  $\mu g/$  0.25 ml agarose in 1 ml, for IP.

### **APPLICATIONS**

Lamin A/C (346) is recommended for detection of Lamin A and Lamin C of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for Lamin A/C siRNA (h): sc-35776, Lamin A/C siRNA (m): sc-29385, Lamin A/C shRNA Plasmid (h): sc-35776-SH, Lamin A/C shRNA Plasmid (m): sc-29385-SH, Lamin A/C shRNA (h) Lentiviral Particles: sc-35776-V and Lamin A/C shRNA (m) Lentiviral Particles: sc-29385-V.

Molecular Weight of Lamin A/C: 69/62 kDa.

Positive Controls: U-251-MG whole cell lysate: sc-364176, C6 whole cell lysate: sc-364373 or HeLa whole cell lysate: sc-2200.

#### **STORAGE**

Store at 4° C, \*\*D0 NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

# **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

#### DATA





Lamin A/C (346): sc-7293. Western blot analysis of Lamin A/C expression in U-251-MG (A), C6 (B), HeLa (C) and NIH/3T3 (D) whole cell lysates.

Lamin A/C (346): sc-7293. Immunofluorescence staining of methanol-fixed Hs68 cells showing nuclear envelope staining.

## SELECT PRODUCT CITATIONS

- dos Santos, O.F., et al. 1991. Nephrotoxicity of cyclosporine: the role of platelet-activating factor and thromboxane. Lipids 26: 1320-1323.
- Ho, J.J., et al. 2012. Functional importance of dicer protein in the adaptive cellular response to hypoxia. J. Biol. Chem. 287: 29003-29020.
- Ward, T.M., et al. 2013. Truncated p110 ERBB2 induces mammary epithelial cell migration, invasion and orthotopic xenograft formation, and is associated with loss of phosphorylated Stat5. Oncogene 32: 2463-2474.
- Xu, Y., et al. 2014. U12, a UDCA derivative, acts as an anti-hepatoma drug lead and inhibits the mTOR/S6K1 and cyclin/CDK complex pathways. PLoS ONE 9: e113479.
- 5. Marchi, S., et al. 2015. Defective autophagy is a key feature of cerebral cavernous malformations. EMBO Mol. Med. 7: 1403-1417.
- Ocampo, A., et al. 2016. *In vivo* amelioration of age-associated hallmarks by partial reprogramming. Cell 167: 1719-1733.e12.
- Zhao, Y., et al. 2017. Small molecule p300/catenin antagonist enhances hematopoietic recovery after radiation. PLoS ONE 12: e0177245.
- Chi, B., et al. 2018. The neurodegenerative diseases ALS and SMA are linked at the molecular level via the ASC-1 complex. Nucleic Acids Res. 46: 11939-11951.
- Mao, G., et al. 2019. Regulation of PLPP3 gene expression by NFκB family transcription factors. J. Biol. Chem. 294: 14009-14019.
- Alqahtani, T., et al. 2020. Salinomycin and its derivatives as potent RET transcriptional inhibitors for the treatment of medullary thyroid carcinoma. Int. J. Oncol. 56: 348-358.



See Lamin A/C (E-1): sc-376248 for Lamin A/C antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor<sup>®</sup> 488, 546, 594, 647, 680 and 790.