SANTA CRUZ BIOTECHNOLOGY, INC.

eotaxin-2 (61016.11): sc-73534



BACKGROUND

Chemokines have been implicated in the regulation of stem/progenitor cell proliferation and movement. The C-C chemokine eotaxin-2 (also known as MPIF-2, CK β -6, or small inducible cytokine A24), which promotes chemotaxis and Ca²⁺ mobilization in human eosinophils, exerts its activity solely through the CCR3 receptor. In addition, eotaxin-2 lacks suppressive activity against immature subsets of myeloid progenitors, which have been stimulated to proliferate by multiple growth factors. While eotaxin-2 is functionally similar to eotaxin, they share only 39% amino acid homology and differ almost completely in the NH2-terminal region.

REFERENCES

- Patel, V.P., et al. 1997. Molecular and functional characterization of two novel human C-C chemokines as inhibitors of two distinct classes of myeloid progenitors. J. Exp. Med. 185: 1163-1172.
- Forssmann, U., et al. 1997. Eotaxin-2, a novel C-C chemokine that is selective for the chemokine receptor CCR3, and acts like eotaxin on human eosinophil and basophil leukocytes. J. Exp. Med. 185: 2171-2176.
- White, J.R., et al. 1997. Cloning and functional characterization of a novel human C-C chemokine that binds to the CCR3 receptor and activates human eosinophils. J. Leukoc. Biol. 62: 667-675.
- Elsner, J., et al. 1998. Eotaxin-2 activates chemotaxis-related events and release of reactive oxygen species via pertussis toxin-sensitive G proteins in human eosinophils. Eur. J. Immunol. 28: 2152-2158.
- Broxmeyer, H.E., et al. 1999. Effects of C-C, C-X-C, C, and CX3C chemokines on proliferation of myeloid progenitor cells, and insights into SDF-1-induced chemotaxis of progenitors. Ann. N.Y. Acad. Sci. 872: 142-162.
- Kitaura, M., et al. 1999. Molecular cloning of a novel human C-C chemokine (eotaxin-3) that is a functional ligand of CC chemokine receptor 3. J. Biol. Chem. 274: 27975-27980.

CHROMOSOMAL LOCATION

Genetic locus: CCL24 (human) mapping to 7q11.23.

SOURCE

eotaxin-2 (61016.11) is a mouse monoclonal antibody raised against recombinant eotaxin-2 of human origin.

PRODUCT

Each vial contains 100 $\mu g~lgG_1$ in 1.0 ml PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, **D0 NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

APPLICATIONS

eotaxin-2 (61016.11) is recommended for detection of eotaxin-2 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000).

Suitable for use as control antibody for eotaxin-2 siRNA (h): sc-63311, eotaxin-2 shRNA Plasmid (h): sc-63311-SH and eotaxin-2 shRNA (h) Lentiviral Particles: sc-63311-V.

Molecular Weight of eotaxin-2: 10 kDa.

RESEARCH USE

For research use only, not for use in diagnostic procedures.