

Myeloid Marker (818-18): sc-73542

BACKGROUND

Myeloid cells originate in the bone marrow during hematopoiesis and encompass all hemopoietic cells except the lymphoid cells (T cells, B cells, NK cells and dendritic cells). Vascular endothelial cells can differentiate from common myeloid progenitors, and these cells that form the bone marrow-derived myeloid lineage express markers such as CD31, von Willebrand factor and Tie2. Other myeloid markers may be used to track certain diseases, such as Kawasaki disease, a self-limited vasculitis that affects many organs, including the skin and mucous membranes, lymph nodes, blood vessel walls and heart.

REFERENCES

1. Bettelheim, P., Paietta, E., Majdic, O., Gadner, H., Schwarzmeier, J. and Knapp, W. 1982. Expression of a myeloid marker on TdT⁺ acute lymphocytic leukemic cells: evidence by double-fluorescence staining. *Blood* 60: 1392-1396.
2. Dolinsky, Z.S., Morse, D.E., Kaplan, R.F., Meyer, R.E., Corry, D. and Pomerleau, O.F. 1987. Neuroendocrine, psychophysiological and subjective reactivity to an alcohol placebo in male alcoholic patients. *Alcohol. Clin. Exp. Res.* 11: 296-300.
3. Gabius, S., Joshi, S.S., Gabius, H.J. and Sharp, J.G. 1991. Establishment, characterization and determination of cell surface sugar receptor (lectin) expression by neoglycoenzymes of a human myeloid marker-expressing B lymphoblastoid cell line. *Anticancer Res.* 11: 793-800.
4. Welker, P., Grabbe, J., Zuberbier, T., Guhl, S. and Henz, B.M. 2000. Mast cell and myeloid marker expression during early *in vitro* mast cell differentiation from human peripheral blood mononuclear cells. *J. Invest. Dermatol.* 114: 44-50.
5. Fujiyama, S., Amano, K., Uehira, K., Yoshida, M., Nishiwaki, Y., Nozawa, Y., Jin, D., Takai, S., Miyazaki, M., Egashira, K., Imada, T., Iwasaka, T. and Matsubara, H. 2003. Bone marrow monocyte lineage cells adhere on injured endothelium in a monocyte chemoattractant protein-1-dependent manner and accelerate reendothelialization as endothelial progenitor cells. *Circ. Res.* 93: 980-989.
6. Lin, Y., Roberts, T.J., Sriram, V., Cho, S. and Brutkiewicz, R.R. 2003. Myeloid marker expression on antiviral CD8⁺ T cells following an acute virus infection. *Eur. J. Immunol.* 33: 2736-2743.
7. Viemann, D., Strey, A., Janning, A., Jurk, K., Klimmek, K., Vogl, T., Hirono, K., Ichida, F., Foell, D., Kehrel, B., Gerke, V., Sorg, C. and Roth, J. 2005. Myeloid-related proteins 8 and 14 induce a specific inflammatory response in human microvascular endothelial cells. *Blood* 105: 2955-2962.
8. Bailey, A.S., Willenbring, H., Jiang, S., Anderson, D.A., Schroeder, D.A., Wong, M.H., Grompe, M. and Fleming, W.H. 2006. Myeloid lineage progenitors give rise to vascular endothelium. *Proc. Natl. Acad. Sci. USA* 103: 13156-13161.
9. Frey, A.B. 2006. Myeloid suppressor cells regulate the adaptive immune response to cancer. *J. Clin. Invest.* 116: 2587-2590.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

SOURCE

Myeloid Marker (818-18) is a mouse monoclonal antibody raised against peripheral blood mononuclear cells of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

Myeloid Marker (818-18) is recommended for detection of myeloid specific DNA binding protein in myeloid precursor cells and leukemias derived from them of human origin by immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for Myeloid Marker siRNA (h): sc-72128, Myeloid Marker shRNA Plasmid (h): sc-72128-SH and Myeloid Marker shRNA (h) Lentiviral Particles: sc-72128-V.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850. 2) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.