

GABA_A Rα5 (N-19): sc-7357

BACKGROUND

GAD-65 and GAD-67, glutamate decarboxylases function to catalyze the production of GABA (γ-aminobutyric acid). In the central nervous system GABA functions as the main inhibitory transmitter by increasing a Cl⁻ conductance that inhibits neuronal firing. GABA has been shown to activate both ionotropic (GABA_A) and metabotropic (GABA_B) receptors as well as a third class of receptors called GABA_C. Both GABA_A and GABA_C are ligand-gated ion channels, however, they are structurally and functionally distinct. Members of the GABA_A receptor family include GABA_A Rα1-6, GABA_A Rβ1-3, GABA_A Rγ1-3, GABA_A Rδ, GABA_A Rε, GABA_A Rρ1 and GABA_A Rρ2. The GABA_B family is composed of GABA_B R1α and GABA_B R1β. GABA transporters have also been identified and include GABA T-1, GABA T-2 and GABA T-3 (also designated GAT-1, -2, and -3). The GABA transporters function to terminate GABA action.

CHROMOSOMAL LOCATION

Genetic locus: GABRA5 (human) mapping to 15q12; Gabra5 (mouse) mapping to 7 C.

SOURCE

GABA_A Rα5 (N-19) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of GABA_A Rα5 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-7357 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

GABA_A Rα5 (N-19) is recommended for detection of GABA_A Rα5 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

GABA_A Rα5 (N-19) is also recommended for detection of GABA_A Rα5 in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for GABA_A Rα5 siRNA (h): sc-42433, GABA_A Rα5 siRNA (m): sc-42434, GABA_A Rα5 siRNA (r): sc-270130, GABA_A Rα5 shRNA Plasmid (h): sc-42433-SH, GABA_A Rα5 shRNA Plasmid (m): sc-42434-SH, GABA_A Rα5 shRNA Plasmid (r): sc-270130-SH, GABA_A Rα5 shRNA (h) Lentiviral Particles: sc-42433-V, GABA_A Rα5 shRNA (m) Lentiviral Particles: sc-42434-V and GABA_A Rα5 shRNA (r) Lentiviral Particles: sc-270130-V.


Molecular Weight of GABA_A Rα5: 55 kDa.

Positive Controls: GABA_A Rα5 (h): 293T Lysate: sc-111120.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



GABA_A Rα5 (N-19): sc-7357. Western blot analysis of GABA_A Rα5 expression in non-transfected: CHO-K1 (A) and human GABA_A Rα5 transfected: sc-111120 (B) whole cell lysates.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **GABA_A Rα1-6 (E-8): sc-376282** or **GABA_A Rα5 (A-5): sc-393921**, our highly recommended monoclonal alternatives to GABA_A Rα5 (N-19). Also, for AC, HRP, FITC, PE, Alexa Fluor® 488 and Alexa Fluor® 647 conjugates, see **GABA_A Rα1-6 (E-8): sc-376282**.