

# Bcl-2 (C-2): sc-7382



The Power to Question

## BACKGROUND

Bcl-2 is one among many key regulators of apoptosis, which are essential for proper development, tissue homeostasis, and protection against foreign pathogens. Human Bcl-2 is an anti-apoptotic, membrane-associated oncoprotein that can promote cell survival through protein-protein interactions with other Bcl-2 related family members, such as the death suppressors Bcl-x<sub>L</sub>, Mcl-1, Bcl-w, and A1 or the death agonists Bax, Bak, Bik, Bad, and Bid. The anti-apoptotic function of Bcl-2 can also be regulated through proteolytic processing and phosphorylation. Bcl-2 may promote cell survival by interfering with the activation of the cytochrome c/Apaf-1 pathway through stabilization of the mitochondrial membrane. Mutations in the Bcl-2 gene can contribute to cancers where normal physiological cell death mechanisms are compromised by deregulation of the anti-apoptotic influence of Bcl-2.

## CHROMOSOMAL LOCATION

Genetic locus: BCL2 (human) mapping to 18q21.33; Bcl2 (mouse) mapping to 1 E2.1.

## SOURCE

Bcl-2 (C-2) is a mouse monoclonal antibody raised against amino acids 1-205 of Bcl-2 of human origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Bcl-2 (C-2) is available conjugated to agarose (sc-7382 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-7382 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-7382 PE), fluorescein (sc-7382 FITC), Alexa Fluor® 488 (sc-7382 AF488), Alexa Fluor® 594 (sc-7382 AF594) or Alexa Fluor® 647 (sc-7382 AF647), 200 µg/ml, for IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-7382 AF680) or Alexa Fluor® 790 (sc-7382 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

In addition, Bcl-2 (C-2) is available conjugated to biotin (sc-7382 B), 200 µg/ml, for WB, IHC(P) and ELISA; and to either TRITC (sc-7382 TRITC, 200 µg/ml), PerCP (sc-7382 PerCP), PerCP-Cy5.5 (sc-7382 PCPC5) or Alexa Fluor® 405 (sc-7382 AF405), 100 tests in 2 ml, for IF, IHC(P) and FCM.

## APPLICATIONS

Bcl-2 (C-2) is recommended for detection of Bcl-2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and flow cytometry (1 µg per 1 x 10<sup>6</sup> cells).

Suitable for use as control antibody for Bcl-2 siRNA (h): sc-29214, Bcl-2 siRNA (m): sc-29215, Bcl-2 shRNA Plasmid (h): sc-29214-SH, Bcl-2 shRNA Plasmid (m): sc-29215-SH, Bcl-2 shRNA (h) Lentiviral Particles: sc-29214-V and Bcl-2 shRNA (m) Lentiviral Particles: sc-29215-V.


Molecular Weight of Bcl-2: 26 kDa.

Positive Controls: WEHI-231 whole cell lysate: sc-2213.


## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## DATA



Bcl-2 (C-2): sc-7382. Western blot analysis of Bcl-2 expression in Jurkat (A), WEHI-231 (B), NIH/3T3 (C), PC-12 (D) and A-10 (E) whole cell lysates.



Bcl-2 (C-2): sc-7382. Immunoperoxidase staining of formalin-fixed, paraffin-embedded normal human colon tissue showing cytoplasmic staining of lymphoid cells (A). Immunofluorescence staining of methanol-fixed HL-60 cells showing cytoplasmic localization (B).

## SELECT PRODUCT CITATIONS

- Royuela, M., et al. 2000. IL-2, its receptors, and Bcl-2 and Bax genes in normal, hyperblastic and carcinomatous human prostate: immunohistochemical comparative analysis. *Growth Factors* 18: 135-146.
- Shin, S.J., et al. 2017. The traditional Korean herbal medicine Ga-Gam-Nai-Go-Hyan suppresses testosterone-induced benign prostatic hyperplasia by regulating inflammatory responses and apoptosis. *Exp. Ther. Med.* 13: 1025-1031.
- Borges, K.S., et al. 2017. The aurora kinase inhibitor AMG 900 increases apoptosis and induces chemosensitivity to anticancer drugs in the NCI-H295 adrenocortical carcinoma cell line. *Anticancer Drugs* 28: 634-644.
- Rivero-Segura, N.A., et al. 2017. Prolactin-induced neuroprotection against glutamate excitotoxicity is mediated by the reduction of [Ca<sup>2+</sup>]<sub>i</sub> overload and NFκB activation. *PLoS ONE* 12: e0176910.
- Choi, I.Y., et al. 2017. Dexmedetomidine alleviates cerebral ischemia-induced short-term memory impairment by inhibiting the expression of apoptosis-related molecules in the hippocampus of gerbils. *Exp. Ther. Med.* 13: 107-116.
- Basarslan, S.K., et al. 2017. Protective effects of intralipid and caffeic acid phenyl ester (CAPE) on neurotoxicity induced by ethanol in rats. *Turk. Neurosurg.* 27: 66-73.
- Massihnia, D., et al. 2017. Phospho-Akt overexpression is prognostic and can be used to tailor the synergistic interaction of Akt inhibitors with gemcitabine in pancreatic cancer. *J. Hematol. Oncol.* 10: 9.
- Sun, L., et al. 2017. MIR506 induces autophagy-related cell death in pancreatic cancer cells by targeting the STAT3 pathway. *Autophagy* 13: 703-714.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA