

Contactin 1 (COEx-4): sc-73858

BACKGROUND

Changes in synaptic efficacy can mediate crucial processes during learning and memory formation. Accumulating evidence implicates cell adhesion molecules in activity-dependent synaptic modifications associated with paired-pulse facilitation (PPF), long-term potentiation (LTP) and long-term depression (LTD). Among the cell adhesion molecules involved in these processes are the contactins. Contactins are immunoglobulin superfamily members that play a selective role in synaptic plasticity, PPF and LTD, and may regulate cell-cell interactions contributing to synaptic plasticity in conjunction with other synapse targeting molecules, including paranodin and phosphacan. In addition, contactins are essential components that control expression and distribution of Na⁺ channels in neurons, junctional attachment at the paranode, and ultimately the physiology of the myelinated nerve. The human Contactin 1 gene encodes a 1,018 amino acid protein. The human Contactin 3 gene encodes a 646 amino acid plasmacytoma-associated neuronal glycoprotein. The human Contactin 5 gene encodes a 1,100 amino acid neural adhesion molecule. The human Contactin 6 gene encodes a 1,028 amino acid neural adhesion molecule.

REFERENCES

1. Ranscht, B. 1988. Sequence of contactin, a 130 kDa glycoprotein concentrated in areas of interneuronal contact, defines a new member of the immunoglobulin supergene family in the nervous system. *J. Cell Biol.* 107: 1561-1573.
2. Fields, R.D. and Itoh, K. 1996. Neural cell adhesion molecules in activity-dependent development and synaptic plasticity. *Trends Neurosci.* 19: 473-480.
3. Kazarinova-Noyes, K., Malhotra, J.D., McEwen, D.P., Mattei, L.N., Berglund, E.O., Ranscht, B., Levinson, S.R., Schachner, M., Shrager, P., Isom, L.L. and Xiao, Z.C. 2001. Contactin associates with Na⁺ channels and increases their functional expression. *J. Neurosci.* 21: 7517-7525.
4. Boyle, M.E., Berglund, E.O., Murai, K.K., Weber, L., Peles, E. and Ranscht, B. 2001. Contactin orchestrates assembly of the septate-like junctions at the paranode in myelinated peripheral nerve. *Neuron* 30: 385-397.
5. Murai, K.K., Misner, D. and Ranscht, B. 2002. Contactin supports synaptic plasticity associated with hippocampal long-term depression but not potentiation. *Curr. Biol.* 12: 181-190.
6. LocusLink Report (LocusID: 1272). <http://www.ncbi.nlm.nih.gov/LocusLink/>

CHROMOSOMAL LOCATION

Genetic locus: CNTN1 (human) mapping to 12q12.

SOURCE

Contactin 1 (COEx-4) is a mouse monoclonal antibody raised against the extracellular domain of Contactin 1 of human origin.

PRODUCT

Each vial contains 100 µg IgG₁ in 1.0 ml PBS with < 0.1% sodium azide and protein stabilizer.

APPLICATIONS

Contactin 1 (COEx-4) is recommended for detection of Contactin 1 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for Contactin 1 siRNA (h): sc-43695, Contactin 1 shRNA Plasmid (h): sc-43695-SH and Contactin 1 shRNA (h) Lentiviral Particles: sc-43695-V.

Molecular Weight of Contactin 1: 135 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-mouse IgG-HRP: sc-2005 (dilution range: 1:2000-1:32,000) or Cruz Marker™ compatible goat anti-mouse IgG-HRP: sc-2031 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-mouse IgG-FITC: sc-2010 (dilution range: 1:100-1:400) or goat anti-mouse IgG-TR: sc-2781 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 3) Immunohistochemistry: use ImmunoCruz™: sc-2050 or ABC: sc-2017 mouse IgG Staining Systems.

SELECT PRODUCT CITATIONS

1. Liu, P., et al. 2011. VEGF-C promotes the development of esophageal cancer via regulating CNTN-1 expression. *Cytokine* 55: 8-17.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.