

IL-12 p70 (EQ-7): sc-74151

BACKGROUND

The interleukins (ILs) are a broad family of well characterized cytokines, primarily of hematopoietic cell origin. As new cytokines are molecularly characterized, they are assigned an IL number to maintain a standard nomenclature. ILs are secreted by immune cells that regulate a wide range of immune system functions. IL-12 is secreted by macrophages and human B-lymphoblastoid cells in response to antigenic stimulation. IL-12 is responsible for the differentiation of naive CD4⁺ T cells into type 1 helper T cells that produce interferon- γ (IFN- γ). It also activates production of tumor necrosis factor α (TNF α) from T and natural killer (NK) cells, and it inhibits IL-4 mediated suppression of IFN- γ . IL-12 also has anti-angiogenic activity, since the production of IFN- γ increases the production of inducible protein-10 (IP-10). IL-12 p40 represents a subunit of IL-12 that is induced by several pathogens. IL-12 p70 is a subunit of IL-12 that is controlled through both the p35 and p40 genes in a cell type-specific fashion.

REFERENCES

1. Yao, L., et al. 2000. Effective targeting of tumor vasculature by the angiogenesis inhibitors vasostatin and interleukin-12. *Blood* 96: 1900-1905.
2. Muller-Suur, C., et al. 2002. Organic dust-induced interleukin-12 production activates T and natural killer cells. *Eur. Respir. J.* 20: 686-690.
3. Noble, A., et al. 2003. CD8⁺ immunoregulatory cells in the graft-versus-host reaction: CD8 T cells activate dendritic cells to secrete interleukin-12/interleukin-18 and induce T helper 1 autoantibody. *Immunology* 109: 476-486.
4. Yamamoto, N., et al. 2004. Essential role for the p40 subunit of interleukin-12 in neutrophil-mediated early host defense against pulmonary infection with *Streptococcus pneumoniae*: involvement of interferon- γ . *Microbes Infect.* 6: 1241-1249.
5. Puertollano, M.A., et al. 2005. Assessment of interleukin-12, interferon- γ , and tumor necrosis factor α secretion in sera from mice fed with dietary lipids during different stages of *Listeria monocytogenes* infection. *Clin. Diagn. Lab. Immunol.* 12: 1098-1103.
6. Entleutner, M., et al. 2005. Impact of interleukin-12, oxidative burst, and iNOS on the survival of murine fecal peritonitis. *Int. J. Colorectal Dis.* 21: 64-70.
7. Gafa, V., et al. 2006. Human dendritic cells following *Aspergillus fumigatus* infection express the CCR7 receptor and a differential pattern of interleukin-12 (IL-12), IL-23, and IL-27 cytokines, which lead to a Th1 response. *Infect. Immun.* 74: 1480-1489.
8. Rentzos, M., et al. 2006. Interleukin-12 is reduced in cerebrospinal fluid of patients with Alzheimer's disease and frontotemporal dementia. *J. Neurol. Sci.* 249: 110-114.

CHROMOSOMAL LOCATION

Genetic locus: Il12a (mouse) mapping to 3 E1, Il12b (mouse) mapping to 11 B1.1.

SOURCE

IL-12 p70 (EQ-7) is a rat monoclonal antibody raised against full length recombinant IL-12 of mouse origin.

PRODUCT

Each vial contains 100 μ g IgG₁ in 1.0 ml of PBS with < 0.1% sodium azide and protein stabilizer.

APPLICATIONS

IL-12 p70 (EQ-7) is recommended for detection of IL-12 p70 heterodimer of mouse origin by WB (non-reducing), ELISA; non cross-reactive with IL-12 p40.

Molecular Weight of IL-12 p70: 70 kDa.

SELECT PRODUCT CITATIONS

1. Rochet, E., et al. 2019. Type I ROP16 regulates retinal inflammatory responses during ocular toxoplasmosis. *PLoS ONE* 14: e0214310.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.