

# FGF-2 (A-7): sc-74413

## BACKGROUND

Fibroblast growth factor-1 (FGF-1), also designated acidic FGF, and fibroblast growth factor-2 (FGF-2), also referred to as basic FGF, are members of a family of growth factors that stimulate proliferation of cells of mesenchymal, epithelial and neuroectodermal origin. Additional members of the FGF family include the oncogenes FGF-3 (Int-2) and FGF-4 (HST/Kaposi), FGF-5, FGF-6, FGF-7 (KGF), FGF-8 (AIGF), FGF-9 (GAF) and FGF-10. Members of the FGF family share 30-55% amino acid sequence identity and similar gene structure, and are capable of transforming cultured cells when overexpressed in transfected cells. Cellular receptors for FGFs are members of a second multigene family including four tyrosine kinases, designated Flg (FGFR-1), Bek (FGFR-L), TKF and FGFR-3.

## REFERENCES

1. Moore, R., et al. 1986. Sequence, topography and protein coding potential of mouse Int-2: a putative oncogene activated by mouse mammary tumor virus. *EMBO J.* 5: 919-924.
2. Delli Bovi, P., et al. 1987. An oncogene isolated by transfection of Kaposi's sarcoma DNA encodes a growth factor that is a member of the FGF family. *Cell* 50: 729-737.

## CHROMOSOMAL LOCATION

Genetic locus: FGF2 (human) mapping to 4q27; Fgf2 (mouse) mapping to 3 B.

## SOURCE

FGF-2 (A-7) is a mouse monoclonal antibody raised against amino acids 10-140 of FGF-2 of human origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>2a</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## APPLICATIONS

FGF-2 (A-7) is recommended for detection of precursor and mature FGF-2 of mouse, rat, human and *Xenopus laevis* origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for FGF-2 siRNA (h): sc-39446, FGF-2 siRNA (m): sc-39447, FGF-2 siRNA (r): sc-108055, FGF-2 shRNA Plasmid (h): sc-39446-SH, FGF-2 shRNA Plasmid (m): sc-39447-SH, FGF-2 shRNA Plasmid (r): sc-108055-SH, FGF-2 shRNA (h) Lentiviral Particles: sc-39446-V, FGF-2 shRNA (m) Lentiviral Particles: sc-39447-V and FGF-2 shRNA (r) Lentiviral Particles: sc-108055-V.


Molecular Weight of FGF-2 isoforms: 18/21/24 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, WI-38 whole cell lysate: sc-364260 or XP12RO whole cell lysate: sc-364364.


## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## DATA



FGF-2 (A-7): sc-74413. Western blot analysis of FGF-2 expression in WI-38 (A) and XP12RO (B) whole cell lysates.



FGF-2 (A-7): sc-74413. Western blot analysis of human recombinant FGF-2.

## SELECT PRODUCT CITATIONS

1. Upadhyay, A., et al. 2014. *Ixora coccinea* enhances cutaneous wound healing by upregulating the expression of collagen and basic fibroblast growth factor. *ISRN Pharmacol.* 2014: 751824.
2. Andreucci, E., et al. 2016. Roles of different IRES-dependent FGF2 isoforms in the acquisition of the major aggressive features of human metastatic melanoma. *J. Mol. Med.* 95: 97-108.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.



See **FGF-2 (G-2): sc-365106** for FGF-2 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.