

# CKR-2B (H-9): sc-74491

## BACKGROUND

C-C or  $\beta$ -chemokine family members are characterized by a pair of adjacent cysteine residues and serve as potent chemoattractants and activators of monocytes and T cells. C-C chemokine receptor family members include CKR-1, CKR-2A, CKR-2B, CKR-3, CKR-4, CKR-5, CKR-6, CKR-7, CKR-8, CKR-9, CKR-10 and the Duffy blood group antigen. Each of these receptors are G protein-coupled, seven pass transmembrane domain proteins whose major physiological role is to function in the chemotaxis of T cells and phagocytic cells to areas of inflammation. However, this receptor family has also been shown to facilitate viral infection. CKR-2 (C-C chemokine receptor type 2) is a 374 amino acid multi-pass membrane protein that belongs to the C-C chemokine receptor family and is expressed as two isoforms, designated CKR-2A and CKR-2B. Both CKR-2 isoforms function as receptors for a variety of proteins, including MCP-1 and MCP-3, thereby influencing intracellular calcium levels and affecting signal transduction throughout the cell.

## REFERENCES

- Schweickart, V.L., et al. 1994. Cloning of human and mouse EBI1, a lymphoid-specific G protein-coupled receptor encoded on human chromosome 17q12-q21.2. *Genomics* 23: 643-650.
- Deng, H., et al. 1996. Identification of a major co-receptor for primary isolates of HIV-1. *Nature* 381: 661-666.
- Dragic, T., et al. 1996. HIV-1 entry into CD4<sup>+</sup> cells is mediated by the chemokine receptor C-C-CKR-5. *Nature* 381: 667-673.
- Feng, Y., et al. 1996. HIV-1 entry cofactor: functional cDNA cloning of a seven-transmembrane, G protein-coupled receptor. *Science* 272: 872-877.
- Alkhatib, G., et al. 1996. C-C CKR-5: a RANTES, MIP-1 $\alpha$ , MIP-1 $\beta$  receptor as a fusion cofactor for macrophage-tropic HIV-1. *Science* 272: 1955-1958.
- Choe, H., et al. 1996. The  $\beta$ -chemokine receptors CCR-3 and CCR-5 facilitate infection by primary HIV-1 isolates. *Cell* 85: 1135-1148.
- Doranz, B.J., et al. 1996. A dual-tropic primary HIV-1 isolate that uses fusin and the  $\beta$ -chemokine receptors CKR-5, CKR-3 and CKR-2B as fusion cofactors. *Cell* 85: 1149-1158.
- Baba, M., et al. 1997. Identification of CCR-6, the specific receptor for a novel lymphocyte-directed C-C chemokine LARC. *J. Biol. Chem.* 272: 14893-14898.

## CHROMOSOMAL LOCATION

Genetic locus: FLJ78302 (human) mapping to 3p21.31.

## SOURCE

CKR-2B (H-9) is a mouse monoclonal antibody raised against amino acids 1-40 mapping at the N-terminus of CKR-2B of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## APPLICATIONS

CKR-2B (H-9) is recommended for detection of CKR-2B of human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for CKR-2B siRNA (h): sc-39883, CKR-2B shRNA Plasmid (h): sc-39883-SH and CKR-2B shRNA (h) Lentiviral Particles: sc-39883-V.

Molecular Weight of CKR-2B: 41 kDa.


Positive Controls: Hep G2 cell lysate: sc-2227.

## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:

- Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## DATA



CKR-2B (H-9): sc-74491. Western blot analysis of human recombinant CKR-2B fusion protein.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.