


## c-Jun (G-4): sc-74543



The Power to Question

## BACKGROUND

Genes belonging to the Jun and Fos oncogene families encode nuclear proteins that are found to be associated with a number of transcriptional complexes. The c-Jun protein is a major component of the transcription factor AP-1, originally shown to mediate phorbol ester tumor promoter (TPA)-induced expression of responsive genes through the TPA response element (TRE). The Jun proteins form homo- and heterodimers which bind the TRE, while Fos proteins are active only as heterodimers with any of the Jun proteins. Fos/Jun heterodimers have a much higher affinity for the TRE than Jun homodimers. Ha-Ras augments c-Jun activity and stimulates phosphorylation of its activation domain. An inhibitor of Fos/Jun function, termed IP-1, associates with Fos and Jun and is inactivated upon phosphorylation induced by the cAMP-dependent protein kinase A (PKA).

## REFERENCES

1. Sambucetti, L.C., et al. 1986. The Fos protein complex is associated with DNA in isolated nuclei and binds to DNA cellulose. *Science* 234: 1417-1419.
2. Bohmann, D., et al. 1987. Human proto-oncogene c-Jun encodes a DNA binding protein with structural and functional properties of transcription factor AP-1. *Science* 238: 1386-1392.
3. Distel, R.J., et al. 1987. Nucleoprotein complexes that regulate gene expression in adipocyte differentiation: direct participation of c-Fos. *Cell* 49: 835-844.

## CHROMOSOMAL LOCATION

Genetic locus: JUN (human) mapping to 1p32.1; Jun (mouse) mapping to 4 C5.

## SOURCE

c-Jun (G-4) is a mouse monoclonal antibody raised against amino acids 1-79 of c-Jun of human origin.

## PRODUCT

Each vial contains 200 µg IgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-74543 X, 200 µg/0.1 ml.

c-Jun (G-4) is available conjugated to agarose (sc-74543 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-74543 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-74543 PE), fluorescein (sc-74543 FITC), Alexa Fluor® 488 (sc-74543 AF488), Alexa Fluor® 546 (sc-74543 AF546), Alexa Fluor® 594 (sc-74543 AF594) or Alexa Fluor® 647 (sc-74543 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-74543 AF680) or Alexa Fluor® 790 (sc-74543 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

c-Jun (G-4) is recommended for detection of c-Jun p39 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Suitable for use as control antibody for c-Jun siRNA (h): sc-29223, c-Jun siRNA (m): sc-29224, c-Jun siRNA (r): sc-156028, c-Jun shRNA Plasmid (h): sc-29223-SH, c-Jun shRNA Plasmid (m): sc-29224-SH, c-Jun shRNA Plasmid (r): sc-156028-SH, c-Jun shRNA (h) Lentiviral Particles: sc-29223-V, c-Jun shRNA (m) Lentiviral Particles: sc-29224-V and c-Jun shRNA (r) Lentiviral Particles: sc-156028-V.

c-Jun (G-4) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of c-Jun: 39 kDa.

Positive Controls: BYDP whole cell lysate: sc-364368, NIH/3T3 whole cell lysate: sc-2210 or NIH/3T3 + PMA nuclear extract: sc-2125.

## DATA



c-Jun (G-4) HRP: sc-74543 HRP. Direct western blot analysis of c-Jun expression in HeLa + UV (A), NIH/3T3 (B) and BYDP (C) whole cell lysates.



c-Jun (G-4): sc-74543. Immunofluorescence staining of formalin-fixed Hep G2 cells showing nuclear localization (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human ovary tissue showing nuclear staining of ovarian stroma cells (B).

## SELECT PRODUCT CITATIONS

1. Wang, Y.H., et al. 2007. IL-25 augments type 2 immune responses by enhancing the expansion and functions of TSLP-DC-activated Th2 memory cells. *J. Exp. Med.* 204: 1837-1847.
2. Qu, Y., et al. 2018. Nobiletin prevents cadmium-induced neuronal apoptosis by inhibiting reactive oxygen species and modulating JNK/ERK1/2 and Akt/mTOR networks in rats. *Neurol. Res.* 40: 211-220.
3. Yan, F., et al. 2018. ATF3 is positively involved in particulate matter-induced airway inflammation *in vitro* and *in vivo*. *Toxicol. Lett.* 287: 113-121.
4. He, H., et al. 2018. c-Jun/AP-1 overexpression reprograms ERα signaling related to tamoxifen response in ERα-positive breast cancer. *Oncogene*. E-published.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.